

ACITRETIN

Products Affected

- Acitretin

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Severely impaired liver or kidney function. Chronic abnormally elevated blood lipid values. Concomitant use of methotrexate or tetracyclines. Pregnancy. Females of child-bearing potential who intend to become pregnant during therapy or at any time for at least 3 years after discontinuing therapy. Females of child-bearing potential who will not use reliable contraception while undergoing treatment and for at least 3 years following discontinuation. Females of child-bearing potential who drink alcohol during treatment or for two months after cessation of therapy.
Required Medical Information	Diagnosis of plaque psoriasis and documented treatment failure, intolerance, or contraindication to any one of the following: high potency steroids, (i.e. betamethasone, fluocinonide, desoximetasone), calcipotriene, or tazarotene.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist
Coverage Duration	12 months
Other Criteria	None

ACTEMRA

Products Affected

- Actemra

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patients with an ANC less than 2000/mm ³ , a platelet count less than 100,000/mm ³ , or an ALT or AST greater than 1.5 times the upper limit of normal. Patient is not receiving Actemra in combination with a biologic DMARD (Enbrel , Humira , Cimzia , Simponi) . Patient is not receiving Actemra in combination with a Janus kinase inhibitor (eg, Xeljanz).
Required Medical Information	Diagnosis of Chimeric antigen receptor (CAR) T cell-induced severe or life-threatening cytokine release syndrome (CRS), Giant cell arteritis, Polyarticular juvenile idiopathic arthritis, Rheumatoid arthritis, OR systemic juvenile idiopathic arthritis. For PAJIA, member needs trial or intolerance/contraindication to Humira. For RA, member needs trial or intolerance/contraindication to Humira and Enbrel. For diagnosis of systemic juvenile idiopathic arthritis, chimeric antigen receptor T cell-induced severe or life-threatening cytokine release syndrome, or Giant cell arteritis Actemra will be approved.
Age Restrictions	2 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ADAGEN

Products Affected

- Adagen

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Severe thrombocytopenia
Required Medical Information	Diagnosis of adenosine deaminase (ADA) deficiency in a patient with severe combined immunodeficiency disease (SCID) AND patient is not a suitable candidate for, or who has failed, bone marrow transplantation.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ADCIRCA

Products Affected

- Adcirca

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patient requires nitrate therapy on a regular or intermittent basis.
Required Medical Information	PAH (WHO Group 1) was confirmed by right heart catheterization.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ADEMPAS

Products Affected

- Adempas

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form. Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline). Pregnancy.
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO group I) AND diagnosis was confirmed by right heart catheterization OR Patient has a diagnosis of chronic thromboembolic pulmonary hypertension (CTEPH, WHO group 4) AND patient has persistent or recurrent disease after surgical treatment (e.g., pulmonary endarterectomy) or has CTEPH that is inoperable AND female patients are enrolled in the ADEMPAS REMS program.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	6 months - initial. 12 months - renewal
Other Criteria	For renewal, medication was effective (i.e. improved 6 minute walk distance, oxygen saturation, etc.)

AFINITOR

Products Affected

- Afinitor

- Afinitor Disperz

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced metastatic renal cell carcinoma and patient has failed therapy (disease progressed) with sunitinib or sorafenib OR in combination with lenvatinib, following one prior anti-angiogenic therapy. Diagnosis of pancreatic neuroendocrine tumors (pNET) that are unresectable, locally advanced or metastatic OR Diagnosis of renal angiomyolipoma with tuberous sclerosis complex (TSC) and patient does not require immediate surgery OR Diagnosis of advanced hormone receptor-positive, HER2-negative breast cancer and patient is a postmenopausal woman and patient has failed treatment with Femara or Arimidex and the medication will be used in combination with Aromasin OR Diagnosis of subependymal giant cell astrocytoma (SEGA) associated with TSC that requires therapeutic intervention but is not a candidate for curative surgical resection OR progressive, well-differentiated, non-functional, neuroendocrine tumors (NET) of gastrointestinal (GI) or lung origin with unresectable, locally advanced or metastatic disease OR renal angiomyolipoma and tuberous sclerosis complex not requiring immediate surgery OR Diagnosis of tuberous sclerosis complex (TSC)-associated partial onset seizures.
Age Restrictions	18 years of age or older for RCC, pNET, and renal angiomyolipoma with TSC. 1 year of age or older for SEGA. 2 years of age or older for TSC associated partial seizures.
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None

ALDURAZYME

Products Affected

- Aldurazyme

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Hurler or Hurler-Scheie form of Mucopolysaccharidosis I (MPS I) or Diagnosis of Scheie form of MPS I with moderate to severe symptoms.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ALECENSA

Products Affected

- Alecensa

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic anaplastic lymphoma kinase positive non-small cell lung cancer. Documentation of intolerance or disease progression following therapy with crizotinib
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ALIQOPA

Products Affected

- Aliqopa

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Follicular Lymphoma: Diagnosis of relapsed follicular lymphoma in patients who have received at least 2 prior systemic therapies
Age Restrictions	18 years of age and older
Prescriber Restrictions	Must be prescribed by, or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of therapy

ALPHA1-PROTEINASE INHIBITOR

Products Affected

- Prolastin-C Intravenous Solution Reconstituted 1000 MG
- Zemaira

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Not covered for patients with IgA deficiency
Required Medical Information	All of the following: A) Patient has an alpha-1 proteinase inhibitor (alpha-1 antitrypsin) deficiency AND B) Diagnosis of emphysema AND C) One of the following: 1) Patient has a high risk phenotype: PiZZ, PiZ(null), Pi(null)(null) OR 2) Patient has serum alpha-1 antitrypsin concentrations of less than 11 uM/L (80 mg/dL) AND D) One of the following: FEV1 level is between 30% and 65% of predicted OR the patient has experienced a rapid decline in lung function (i.e., reduction of FEV1 more than 120 mL/year) that warrants treatment.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Pulmonologist
Coverage Duration	12 months
Other Criteria	None

ALUNBRIG

Products Affected

- Alunbrig

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic, ALK positive non-small cell lung cancer and have progressed or are intolerant to crizotinib.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a oncologist
Coverage Duration	12 months
Other Criteria	None

AMPYRA

Products Affected

- Ampyra

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	History of seizure. Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute).
Required Medical Information	Diagnosis of multiple sclerosis. Patient must demonstrate sustained walking impairment, but with the ability to walk 25 feet (with or without assistance) prior to starting Ampyra and patient is currently on a disease modifying drug (interferon beta 1a, peginterferon beta 1a, intereron beta 1b, or glatiramer) to control disease progression, or has documented treatment failure, intolerance, or contraindication to any one of the following: interferon beta 1a, peginterferon beta 1a, intereron beta 1b, or glatiramer.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Initial - 3 months. Renewal - 12 months
Other Criteria	Multiple Sclerosis (MS) (initial): Diagnosis of MS. Physician confirmation that patient has difficulty walking (eg, timed 25 foot walk test). One of the following: expanded disability status scale (EDSS) score less than or equal to 7, or not restricted to using a wheelchair (if EDSS is not measured).

APOKYN

Products Affected

- Apokyn Subcutaneous Solution Cartridge

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	PD (Initial, reauth): Patient is using Apokyn with any 5-HT3 antagonist (eg, ondansetron, granisetron, dolasetron, palonosetron, alosetron)
Required Medical Information	Parkinson's disease (PD) (Initial): Diagnosis of advanced PD. Patient is experiencing acute intermittent hypomobility (defined as off episodes characterized by muscle stiffness, slow movements, or difficulty starting movements). Patient is receiving Apokyn in combination with other medications for the treatment of PD (e.g., carbidopa/levodopa, pramipexole, ropinirole, benztropine, etc.).
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ARCALYST

Products Affected

- Arcalyst

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of CAPS, including Familial Cold Auto-inflammatory Syndrome (FCAS) and/or Muckle-Wells Syndrome (MWS). The medication will not be used in combination with another biologic.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with or recommendation of, an immunologist, allergist, dermatologist, rheumatologist, neurologist, or other medical specialist
Coverage Duration	12 months
Other Criteria	CAPS (Reauth): Patient has experienced disease stability or improvement in clinical symptoms while on therapy as evidence by one of the following: A) improvement in rash, fever, joint pain, headache, conjunctivitis, B) decreased number of disease flare days, C) normalization of inflammatory markers (CRP, ESR, SAA), D) corticosteroid dose reduction, OR E) improvement in MD global score or active joint count.

ARMODAFINIL

Products Affected

- Armodafinil

- Modafinil

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) excessive sleepiness associated with obstructive sleep apnea (OSA)/hypopnea syndrome confirmed by sleep lab evaluation, e.g., multiple sleep latency test, polysomnography), B) excessive sleepiness associated with narcolepsy confirmed by sleep lab evaluation and patient has tried and failed, is unable to tolerate, or has contraindication(s) to at least one other central nervous system stimulant (e.g., methylphenidate, mixed amphetamine salts, dextroamphetamine), OR C) excessive sleepiness associated with shift work disorder with a primary complaint of excessive sleepiness or insomnia which is temporally associated with a work period (usually night work) that occurs during the habitual sleep phase or polysomnography and the MSLT demonstrate loss of a normal sleep-wake pattern.
Age Restrictions	17 years of age and older
Prescriber Restrictions	None
Coverage Duration	OSA/hypopnea syndrome: 6 months (initial), 12 months (renewal). Other diagnoses: 12 months.
Other Criteria	None

AUBAGIO

Products Affected

- Aubagio

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Severe hepatic impairment. Current treatment with leflunomide. Patients who are pregnant or women of childbearing potential not using reliable contraception.
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis (e.g., relapsing-remitting MS or progressive-relapsing MS) OR patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient has experienced an objective response to therapy (i.e. no or slowed progression of disease)

AURYXIA

Products Affected

- Auryxia

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	For the management of hyperphosphatemia in patients with chronic kidney disease on dialysis
Age Restrictions	18 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

AUSTEDO

Products Affected

- Austedo

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Suicidal ideation and/or untreated or inadequately treated depression, hepatic impairment, or taking MAOIs, reserpine, or tetrabenazine.
Required Medical Information	Diagnosis of Chorea associated with Huntington's disease OR Diagnosis of Tardive dyskinesia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in collaboration with a neurologist or psychiatrist
Coverage Duration	12 months
Other Criteria	Dosing will be approved per the FDA labeling based on CYP2D6 testing. For renewal, patient had an objective response to therapy.

AVASTIN

Products Affected

- Avastin

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Gastrointestinal perforation. Wound dehiscence. Serious hemorrhage or recent hemoptysis.
Required Medical Information	Treatment of: 1) metastatic colorectal cancer: a) with irinotecan, leucovorin, and 5-fluorouracil (5-FU), b) with leucovorin, 5-fluorouracil (5-FU), and oxaliplatin, c) second-line treatment in patients who progressed on a first-line bevacizumab-containing regimen, with fluoropyrimidine and irinotecan or fluoropyrimidine and oxaliplatin-based chemotherapy, d) first-line or second-line treatment with 5-fluorouracil, leucovorin, and irinotecan (IFL plus bevacizumab) OR 2) first-line treatment of unresectable, locally advanced, recurrent, or metastatic non-squamous non-small cell lung cancer (NSCLC) with carboplatin and paclitaxel OR 3) metastatic renal cell carcinoma with interferon alfa OR 4) monotherapy treatment of glioblastoma that has progressed on prior therapy OR 5) ovarian cancer: a) recurrent, platinum-sensitive, epithelial ovarian, fallopian tube, or primary peritoneal cancer, with carboplatin and paclitaxel, b) recurrent, platinum-sensitive, epithelial ovarian, fallopian tube, or primary peritoneal cancer with gemcitabine and carboplatin, c) recurrent, platinum-resistant, epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who have received no more than 2 prior chemotherapy regimens, with paclitaxel, d) recurrent, platinum-resistant, epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who have received no more than 2 prior chemotherapy regimens, with peg-liposomal doxorubicin, e) recurrent, platinum-resistant, epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who have received no more than 2 prior chemotherapy regimens, with topotecan OR 6) cervical cancer: a) persistent, recurrent, or metastatic cervical cancer, with paclitaxel and cisplatin, b) persistent, recurrent, or metastatic cervical cancer, with paclitaxel and topotecan.
Age Restrictions	18 years of age and older

PA Criteria	Criteria Details
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

BAVENCIO

Products Affected

- Bavencio

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic Merkel cell carcinoma or locally advanced or metastatic urothelial carcinoma, in patients with disease progression on or following platinum-containing chemotherapy, or within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	None

BELEODAQ

Products Affected

- Beleodaq

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory peripheral T-cell lymphoma (PTCL) (ANC should be greater than or equal to 1000/mm ³ and platelets should be greater than or equal to 50,000/mm ³ prior to each cycle)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

BENLYSTA

Products Affected

- Benlysta

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Systemic lupus erythematosus (SLE) (init): Diagnosis of active SLE. Autoantibody positive (ie, anti-nuclear antibody [ANA] titer greater than or equal to 1:80 or anti-dsDNA level greater than or equal to 30 IU/mL). Currently receiving at least one standard of care treatment for active SLE (eg, antimalarials [eg, Plaquenil (hydroxychloroquine)], corticosteroids [eg, prednisone], or immunosuppressants [eg, methotrexate, Imuran (azathioprine), CellCept (mycophenolate mofetil)]).
Age Restrictions	None
Prescriber Restrictions	SLE (init): Prescribed by or in consultation with a rheumatologist
Coverage Duration	SLE (init, reauth): 6 months
Other Criteria	SLE (reauth): Documentation of positive clinical response to Benlysta therapy

BEXAROTENE

Products Affected

- Bexarotene

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma (CTCL) and patient is not a candidate for or had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) for cutaneous manifestations of CTCL
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	Female patients of child-bearing potential have a documented negative pregnancy test one week prior to the initiation of therapy. For renewal, Patient has not had disease progression while on therapy and female patients of child-bearing potential are not pregnant and are continuing to use adequate birth-control measures during therapy.

BOSULIF

Products Affected

- Bosulif

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Signed statement of diagnosis from the physician, hepatic panel and CBC, trial and failure of imatinib or dasatinib and documentation of a 90 day response OR newly diagnosed chronic phase Philadelphia chromosome-positive chronic myelogenous leukemia (Ph + CML)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

BRIVIACT

Products Affected

- Briviact

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of partial-onset seizures, member must have history of inadequate response, contraindication, or intolerance to levetiracetam prior to approval.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

BUPRENORPHINE SL

Products Affected

- Buprenorphine HCl Sublingual Tablet
Sublingual 2 MG, 8 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Patient has a diagnosis of opioid dependence
Age Restrictions	16 years of age or older
Prescriber Restrictions	None
Coverage Duration	Initial - 3 months. Renewal - 9 months
Other Criteria	For renewal, patient meets all initial criteria

CABOMETYX

Products Affected

- Cabometyx

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patients who have or are at risk for severe hemorrhage and/or patients with a recent history of bleeding or hemoptysis.
Required Medical Information	Diagnosis of advanced renal cell carcinoma who have received prior anti-angiogenic therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

CALQUENCE

Products Affected

- Calquence

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	MANTLE CELL LYMPHOMA (MCL) (1) Patient must have a diagnosis of MCL AND (2) Patient has tried one other therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Must be prescribed by, or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of therapy

CAPRELSA

Products Affected

- Caprelsa

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Congenital long QT syndrome
Required Medical Information	Diagnosis of symptomatic or progressive medullary thyroid cancer with unresectable locally advanced or metastatic disease
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

CARBAGLU

Products Affected

- Carbaglu

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of N-acetyl glutamate synthase (NAGS) deficiency AND patient is experiencing either acute or chronic hyperammonemia
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

CARIMUNE

Products Affected

- Carimune NF Intravenous Solution Reconstituted 6 GM

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	<p>Diagnosis of a primary humoral immunodeficiency disorder such as: primary immunoglobulin deficiency syndrome X-linked immunodeficiency with hyperimmunoglobulin etc). Documented hypogammaglobulinemia (IgG less than 600mg/dl)</p> <p>Idiopathic/Immune Thrombocytopenia Purpura. Diagnosis of Acute ITP with any of the following: Management of acute bleeding due to severe thrombocytopenia (platelets less than 30 000/uL) To increase platelet counts prior major surgical procedures Severe thrombocytopenia (platelets less than 20 000/uL) at risk for intracerebral hemorrhage. Diagnosis of Chronic ITP and ALL of the following are met: Prior treatment has included corticosteroids and splenectomy Duration of illness less than 6 months No concurrent illness explaining thrombocytopenia Platelets persistently at or below 20 000/uL. Chronic Lymphocytic Leukemia (CLL B-cell). With either of the following present: Hypogammaglobulinemia (IgG less than 600mg/dL) Recurrent bacterial infections associated with B-cell CLL. Kawasaki Disease. Diagnosed with Kawasaki Syndrome within ten days of onset of disease manifestations or is diagnosed after ten days of disease onset and continues to exhibit manifestations of inflammation or evolving coronary artery disease. IVIG is used in combination with high dose aspirin for the prevention of coronary artery aneurysms. Bone Marrow Transplant (BMT). Member is hypogammaglobinemic (IgG less than 400mg/dL). Hematopoietic Stem Cell Transplantation (HSCT). Is within first 100 days of allogeneic hematopoietic stem cell transplantation. Is experiencing hypogammaglobulinemia (serum IgG level less than 400 mg/dL). AIDS/HIV. Has any of the following conditions: CD4+ T-cell counts greater than or equal 200/mm³ To prevent maternal transmission of HIV infection IVIG is used in conjunction with zidovudine to prevent serious bacterial infections in HIV-infected members who have hypogammaglobulinemia (serum IgG less than</p>

PA Criteria	Criteria Details
	400 mg/dL).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

CAYSTON

Products Affected

- Cayston

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis is confirmed by appropriate diagnostic or genetic testing. Confirmation of <i>P. aeruginosa</i> in cultures of the airways. For continuation of therapy, a clinical reason to continue therapy, such as symptomatic improvement or pulmonary function tests have not deteriorated more than 10% from baseline.
Age Restrictions	7 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations)

CIMZIA

Products Affected

- Cimzia Prefilled

- Cimzia Subcutaneous Kit 2 X 200 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Active infection (including TB). Concurrent therapy with other biologics.
Required Medical Information	Screening for latent TB infection and assessment for Hep B risk. For positive latent TB, patient must have completed treatment or is currently receiving treatment for LTBI. HBV infection ruled out or treatment initiated for positive infection. Rheumatoid arthritis (RA) - Must have an inadequate response to one of following: 1) inadequate response to methotrexate, 2) inadequate response to another nonbiologic DMARD (e.g., leflunomide, hydroxychloroquine, sulfasalazine) if contraindicated or intolerant to MTX or, 3) intolerance or contraindication to at least 2 nonbiologic DMARDs. Crohn's Disease - Must have an inadequate response or contraindication/intolerance to at least one oral corticosteroid. For ankylosing spondylitis, psoriatic arthritis, severe plaque psoriasis, and/or rheumatoid arthritis member needs trial or intolerance/contraindication to Humira and Enbrel. For Crohn's, member needs trial or intolerance/contraindication to Humira.
Age Restrictions	18 years of age and older
Prescriber Restrictions	CD (init): Prescribed by or in consultation with a gastroenterologist. RA, AS (init): Prescribed by or in consultation with a rheumatologist. PsA (init): Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	Initial: 3 months for Crohn's disease, 1 year for all other indications. Renewal: Plan Year
Other Criteria	For re-authorization, patient's condition must have improved or stabilized.

CINRYZE

Products Affected

- Cinryze

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	History of life-threatening immediate hypersensitivity reactions, including anaphylaxis to the product.
Required Medical Information	Diagnosis of hereditary angioedema AND Medication will be used for routine prophylaxis against angioedema.
Age Restrictions	None
Prescriber Restrictions	prescribed or overseen by a hematologist or immunologist
Coverage Duration	12 months
Other Criteria	None

COMETRIQ

Products Affected

- Cometriq (100 mg Daily Dose)
- Cometriq (140 mg Daily Dose)
- Cometriq (60 mg Daily Dose)

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Gastrointestinal perforation. Fistula. Severe hemorrhage.
Required Medical Information	Diagnosis of progressive, metastatic medullary thyroid cancer OR Diagnosis of advanced or metastatic renal cell cancer
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

COPAXONE

Products Affected

- Glatiramer Acetate

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsing-remitting multiple sclerosis OR diagnosis of first clinical episode with MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient does not have progressive disease and responding to therapy.

CORLANOR

Products Affected

- Corlanor

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Decompensated acute heart failure, hypotension (i.e. blood pressure less than 90/50 mmHg), sick sinus syndrome, sinoatrial block, or 3rd degree AV block, unless a functioning demand pacemaker is present and bradycardia (i.e., resting heart rate less than 60 bpm prior to treatment)
Required Medical Information	Patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction 35% or less, who are in sinus rhythm with resting heart rate 70 beats per minute or more and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

COSENTYX

Products Affected

- Cosentyx 300 Dose

- Cosentyx Sensoready 300 Dose

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	<p>Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: Failure, contraindication, or intolerance to Enbrel (etanercept) AND Humira (adalimumab), OR for continuation of prior Cosentyx therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), OR for continuation of prior Cosentyx therapy. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), OR for continuation of prior Cosentyx therapy. All indications (Initial, reauth): Patient is not receiving Cosentyx in combination with a biologic DMARD [eg, Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab)]. Patient is not receiving Cosentyx in combination with a Janus kinase inhibitor [eg, Xeljanz (tofacitinib)]. For a diagnosis of PsA or plaque psoriasis, Patient is not receiving Cosentyx in combination with a phosphodiesterase 4 (PDE4) inhibitor [e.g., Otezla (apremilast)].</p>
Age Restrictions	None
Prescriber Restrictions	Plaque psoriasis (Initial): Prescribed by or in consultation with a dermatologist. PsA (Initial): Prescribed by or in consultation with a rheumatologist or dermatologist. AS (Initial): Prescribed by or in consultation with a rheumatologist.
Coverage Duration	All indications (Initial, reauth): 12 months
Other Criteria	All indications (Reauth): Documentation of positive clinical response to Cosentyx therapy.

COTELLIC

Products Affected

- Cotellic

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable OR metastatic malignant melanoma with BRAF V600E OR V600K mutation. Documentation of combination therapy with vemurafenib
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

CYSTARAN

Products Affected

- Cystaran

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Demonstrated cysteamine hypersensitivity or penicillamine hypersensitivity
Required Medical Information	Patient has a diagnosis of cystinosis AND patient has corneal cystine crystal accumulation
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

DALIRESP

Products Affected

- Daliresp

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Moderate to severe liver impairment (Child-Pugh B or C)
Required Medical Information	Diagnosis of severe chronic obstructive pulmonary disease (COPD) (defined as FEV1 less than or equal to 50% of predicted and FEV1/forced vital capacity [FVC] less than 0.7) associated with chronic bronchitis AND history of COPD exacerbations which requires the use of systemic corticosteroids, antibiotics, or hospital admission AND Medication will be used with a long-acting inhaled bronchodilator (i.e. long-acting anticholinergic, or long-acting beta agonist in combination with inhaled corticosteroid) or patient is at high-risk of COPD exacerbation and is not a candidate for long-acting inhaled bronchodilator therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

DARZALEX

Products Affected

- Darzalex Intravenous Solution 100 MG/5ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple myeloma and one of the following: patient has received at least 3 prior lines of therapy including a proteasome inhibitor and an immunomodulatory agent or who are double-refractory to a PI and an immunomodulatory agent or patient has received at least 1 prior therapy, in combination with bortezomib and dexamethasone or patient has received at least 1 prior therapy, in combination with lenalidomide and dexamethasone or patient has received at least 2 prior therapies including lenalidomide and a proteasome inhibitor, in combination with pomalidomide and dexamethasone, or newly diagnosed patient ineligible for autologous stem cell transplant and used in combination with bortezomib, melphalan, and prednisone.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

DICLOFENAC TOPICAL

Products Affected

- Diclofenac Sodium Transdermal Gel

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diclofenac 1%: Diagnosis of osteoarthritis, diclofenac 3% gel: Diagnosis of actinic keratosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ELAPRASE

Products Affected

- Elaprase

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis confirmed by DNA testing or enzymatic analysis (deficiency of iduronate 2-sulfatase enzyme activity).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

EMPLICITI

Products Affected

- Empliciti

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple myeloma, documentation of combination therapy with lenalidomide and dexamethasone. Prescriber must document prior treatment with 1 to 3 previous therapies.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

EMSAM

Products Affected

- Emsam

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pheochromocytoma. Patient is taking or will take any of the following: SSRIs, SNRIs, tricyclic antidepressants (TCAs), bupropion, buspirone, meperidine, tramadol, methadone, pentazocine, dextromethorphan, St. John's wort, mirtazapine, cyclobenzaprine, oral selegiline, other MAOIs, oxcarbazepine, carbamazepine, and/or sympathomimetic amines
Required Medical Information	Diagnosis of major depressive disorder AND Patient had adequate trial with at least 2 generic oral antidepressants from differing classes (at least one should be from the following list: selective serotonin reuptake inhibitors, serotonin and norepinephrine reuptake inhibitors, mirtazapine, or bupropion unless contraindicated), unless unable to take any oral medication AND Patient had an adequate washout period (for patients previously on agents requiring a washout period)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, the patient has improved or stabilized on Emsam.

ENBREL

Products Affected

- Enbrel Subcutaneous Solution Prefilled Syringe

- Enbrel Subcutaneous Solution Reconstituted
- Enbrel SureClick Subcutaneous Solution Auto-Injector

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Active serious infection (including tuberculosis). Combined use with a biologic disease-modifying anti-rheumatic drugs or potent immunosuppressant (e.g., azathioprine or cyclosporine)
Required Medical Information	Diagnosis of moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs for at least 3 consecutive months OR Diagnosis of moderate to severe polyarticular juvenile idiopathic arthritis and patient had an inadequate response, intolerance or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs (DMARDs) for at least 3 consecutive months OR Diagnosis of psoriatic arthritis and patient had an inadequate response, intolerance, or contraindication to methotrexate OR Diagnosis of ankylosing spondylitis and patient had an inadequate response, intolerance or contraindication to one or more NSAIDs OR Diagnosis of moderate to severe chronic plaque psoriasis (affecting more than 5% of body surface area or affecting crucial body areas such as the hands, feet, face, or genitals) and patient had an inadequate response, intolerance or contraindication to conventional therapy with at least one of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least one continuous month or one or more oral systemic treatments (i.e. methotrexate, cyclosporine, acitretin, sulfasalazine) for at least 3 consecutive months.
Age Restrictions	2 years of age or older for JIA or JRA. 4 years of age or older for plaque psoriasis. 18 years of age or older for all other indications
Prescriber Restrictions	None

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	None

ENDARI

Products Affected

- Endari

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute sickle cell disease AND Must have A) trial history of Hydroxyurea OR B) intolerance to Hydroxyurea OR C) contraindication to Hydroxyurea
Age Restrictions	5 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ENTRESTO

Products Affected

- Entresto

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	History of angioedema related to previous ACE inhibitor or ARB therapy, concomitant use or use within 36 hours of ACE inhibitors, concomitant use of aliskiren in patients with diabetes
Required Medical Information	Statement of diagnosis indicating Heart Failure (NYHA Class II through IV)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ERBITUX

Products Affected

- Erbitux Intravenous Solution 100 MG/50ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) Locally or regionally advanced squamous cell carcinoma of the head and neck in combination with radiation therapy B) recurrent locoregional disease or metastatic squamous cell carcinoma of the head and neck in combination with platinum-based therapy with 5-fluorouracil as first-line treatment C) Recurrent or metastatic squamous cell carcinoma of the head and neck for whom prior platinum-based therapy has failed and the medication will be used as a single agent, D) K-Ras mutation-negative (wild-type) epidermal growth factor receptor (EGFR)-expressing, metastatic colorectal cancer as determined by an FDA-approved test and the medication will be used: in combination with FOLFIRI for first-line treatment, in combination with irinotecan in patients who are refractory to irinotecan-based chemotherapy, or as a single agent in patients who have failed oxaliplatin- and irinotecan-based chemotherapy or who are intolerant to irinotecan.
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Subject to B vs D

ERIVEDGE

Products Affected

- Erivedge

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic basal cell carcinoma OR Diagnosis of locally advanced basal cell carcinoma that has recurred following surgery or when the patient is not a candidate for surgery and radiation
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ERLEADA

Products Affected

- Erleada

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of nonmetastatic, castration-resistant prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None

ERWINAZE

Products Affected

- Erwinaze Injection

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ESBRIET

Products Affected

- Esbriet

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Appropriate diagnosis (idiopathic pulmonary fibrosis [IPF]), monitoring (hepatic function/LFTs)
Age Restrictions	None
Prescriber Restrictions	Pulmonologist
Coverage Duration	12 months
Other Criteria	None

ESRD THERAPY

Products Affected

- Aranesp (Albumin Free) Injection Solution 100 MCG/ML, 200 MCG/ML, 25 MCG/ML, 300 MCG/ML, 40 MCG/ML, 60 MCG/ML
- Aranesp (Albumin Free) Injection Solution Prefilled Syringe

- Epogen Injection Solution 10000 UNIT/ML, 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML
- Mircera Injection Solution Prefilled Syringe 100 MCG/0.3ML, 50 MCG/0.3ML, 75 MCG/0.3ML
- Procrit

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Pretreatment hemoglobin levels of less than 10g/dL. Dose reduction or interruption if hemoglobin exceeds 10 g/dL (CKD not on dialysis-adult, cancer), 11 g/dL (CKD on dialysis), 12 g/dL (pediatric CKD) in addition to supporting statement of diagnosis from physician.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None

EXJADE

Products Affected

- Exjade

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Creatinine clearance less than 40 mL/min or evidence of overt proteinuria, platelet count less than 50 x 10 ⁹ /L, advanced malignancy, high-risk myelodysplastic syndrome (MDS) with poor performance status, or concurrent use of deferoxamine or iron-containing products.
Required Medical Information	The patient must meet all of the following criteria: 1) Diagnosis of transfusion-dependent anemia with chronic iron overload due to blood transfusions, 2) Patient will have baseline and monthly monitoring of serum ferritin, serum creatinine, creatinine clearance, serum transaminases, and bilirubin. OR For the treatment of chronic iron overload in patients 10 years and older with nontransfusion-dependent thalassemia syndromes
Age Restrictions	Covered for those 2 years of age and older with chronic iron overload due to blood transfusions
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None

FABRAZYME

Products Affected

- Fabrazyme

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Fabry disease.
Age Restrictions	8 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

FARESTON

Products Affected

- Fareston

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis. Must have previous inadequate response or intolerance to tamoxifen. For reauth: must have chart documentation from prescriber indicating improvement in condition.
Age Restrictions	None
Prescriber Restrictions	Oncologist or hematologist
Coverage Duration	6 months
Other Criteria	None

FARYDAK

Products Affected

- Farydak

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Multiple Myeloma (MM) Used in combination with both of the following: Velcade (bortezomib) and dexamethasone. Patient has received at least two prior treatment regimens which included both of the following: Velcade (bortezomib) and an immunomodulatory agent [eg, Revlimid (lenalidomide), Thalomid (thalidomide)].
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist/hematologist
Coverage Duration	12 months
Other Criteria	None

FENTANYL PATCH

Products Affected

- FentaNYL

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Management of acute or post-operative pain. Opioid non-tolerant patients.
Required Medical Information	Patient is opioid tolerant (taking for one week or longer at least 60mg of morphine or equivalent daily) AND has tried two extended release oral opioids or is unable to take extended release oral opioids secondary to allergy, adverse events, swallowing difficulty, or uncontrollable nausea/vomiting.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

FENTANYL SL

Products Affected

- Abstral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Management of acute or post-operative pain, including headache/migraine, dental pain, or use in the emergency room. Opioid non-tolerant patients.
Required Medical Information	Patient meets the following: A) Diagnosis of cancer and use is for breakthrough cancer pain, B) patient is opioid tolerant and taking at least 60 mg morphine/day, at least 25 mcg transdermal fentanyl/hour, at least 30 mg of oxycodone daily, at least 8 mg oral hydromorphone daily or an equianalgesic dose of another opioid for a week or longer, C) at least one other formulary short-acting strong narcotic analgesic alternatives (other than fentanyl) have been ineffective, not tolerated, or contraindicated, D) prescriber and patient are registered in the Transmucosal Immediate Release Fentanyl (TIRF) Risk Evaluation and Mitigation Strategy Access program, E) for brand requests, generic transmucosal fentanyl citrate has been ineffective or not tolerated.
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

FIRAZYR

Products Affected

- Firazyr

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary angioedema AND medication will be used for the treatment of acute attacks.
Age Restrictions	18 years of age and older
Prescriber Restrictions	prescribed or overseen by a hematologist or immunologist
Coverage Duration	12 months
Other Criteria	None

FIRMAGON

Products Affected

- Firmagon

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

FORTEO

Products Affected

- Forteo Subcutaneous Solution 600 MCG/2.4ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Patient has a diagnosis of one of the following: a) osteoporosis in a postmenopausal female, b) primary or hypogonadal osteoporosis in a male, or c) osteoporosis associated with sustained systemic glucocorticoid therapy AND patient is considered to be at high-risk for fracture by meeting one or more of the following: A) history of osteoporotic fracture, B) Low Bone Density less than 2.5 SD below normal, AND one or more of the following: i) failed one oral bisphosphonate and 1 injectable bisphosphonate, or ii) intolerant to one oral bisphosphonate and one injectable bisphosphonate. Patient has not received more than 2 years of therapy with Forteo.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Approve doses based on FDA labeling

GATTEX

Products Affected

- Gattex

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Active gastrointestinal malignancy (gastrointestinal tract, hepatobiliary, pancreatic), colorectal cancer, or small bowel cancer
Required Medical Information	Diagnosis of short bowel syndrome AND patient is receiving specialized nutritional support (i.e. parenteral nutrition)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient has a reduced need for parenteral support (20% reduction) after at least 6 months of therapy.

GILENYA

Products Affected

- Gilenya Oral Capsule 0.5 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Recent (within the last 6 months) occurrence of: myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III/IV heart failure. History or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome, unless patient has a pacemaker. Baseline QTc interval greater than or equal to 500 ms. Receiving concurrent treatment with Class Ia or Class III anti-arrhythmic drugs (quinidine, procainamide, amiodarone, sotalol).
Required Medical Information	Diagnosis of a relapsing form of multiple sclerosis or diagnosis of first clinical episode with MRI features consistent with MS AND Patient will be observed for signs and symptoms of bradycardia in a controlled setting for at least 6 hours after the first dose
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial - 6 months. Renewal - 12 months
Other Criteria	For renewal, the patient has experienced no or slowed disease progression.

GILOTRIF

Products Affected

- Gilotrif

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from the physician in patients with: 1) metastatic non-small cell lung cancer (NSCLC) whose tumors have non-resistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test or 2) metastatic squamous NSCLC, progressing after platinum-based chemotherapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

GLEOSTINE

Products Affected

- Gleostine Oral Capsule 10 MG, 100 MG, 40 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Statement of diagnosis indicating Hodgkin's disease, OR intracranial tumor, OR carcinoma of the breast, OR colorectal cancer, OR lung cancer, OR malignant melanoma, OR malignant tumor of the thymus, OR multiple myeloma, OR non-Hodgkin's lymphoma. AND monitoring of blood counts for evidence of Bone Marrow Suppression (thrombocytopenia or leukopenia).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

GOCOVRI

Products Affected

- Gocovri

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patients with ESRD (CrCl below 15 ml/min/m ²)
Required Medical Information	INITIAL: Diagnosis of Parkinsons disease AND (1) Patient is experiencing dyskinesia AND (2) Patient is receiving levodopa based therapy AND (3) Must have documented trial and failure to amantadine immediate release. RENEWAL: (1) must meet the initial criteria above AND (2) Documentation of positive clinical response to Gocovri (e.g., decreased "off" periods, decreased "on" time with troublesome dyskinesia)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None

GONADOTROPIN

Products Affected

- Chorionic Gonadotropin Intramuscular

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Fertility indications in females are excluded.
Required Medical Information	Diagnosis of Hypogonadotropic hypogonadism or Prepubertal cryptorchidism
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

GROWTH HORMONE

Products Affected

- Norditropin FlexPro

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HALAVEN

Products Affected

- Halaven

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Breast Cancer: Diagnosis of recurrent or metastatic breast cancer. Previous treatment with both of the following: one anthracycline [eg, doxorubicin, Ellence (epirubicin)] and one taxane [eg, paclitaxel, Taxotere (docetaxel)]. Liposarcoma: Diagnosis of unresectable or metastatic liposarcoma. Previous treatment with one anthracycline-containing regimen.
Age Restrictions	None
Prescriber Restrictions	All Uses: prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

HEPATITIS B

Products Affected

- Adefovir Dipivoxil
- Baraclude Oral Solution
- Entecavir
- Vemlidy

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patients that have immune-tolerant chronic hepatitis B per AASLD guidelines
Required Medical Information	Must submit documentation of immune-active chronic hepatitis B per AASLD guidelines.
Age Restrictions	None
Prescriber Restrictions	Prescriber must be a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	12 months
Other Criteria	None

HEPATITIS C

Products Affected

- Epclusa

- Mavyret
- Zepatier

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Must submit documentation of chronic hepatitis C genotype (confirmed by HCV RNA level within the last 6 months). Must submit laboratory results within 6 weeks of initiating therapy including: 1) CBC w Platelets, 2) AST/ALT, 3) Total Bilirubin, 4) Serum Albumin, 5) PT/INR, 6) Serum Creatinine, and 7) GFR. FOR GENOTYPES 1 and 4: Must include subtype, trail/failure, contraindication to, or intolerance to Zepatier or Mavyret prior to approval of Epclusa
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescriber must be a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	Duration of approval per AASLD Guidelines
Other Criteria	None

HERCEPTIN

Products Affected

- Herceptin

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) HER2 overexpressing breast cancer AND patient is node positive OR node negative and either ER/PR negative or ER/PR positive with one high risk feature (i.e. pathological tumor size greater than 2 cm, Grade 2-3, or age less than 35 years) AND medication is for adjuvant treatment as part of a regimen consisting of: doxorubicin, cyclophosphamide, and either paclitaxel or docetaxel OR with docetaxel and carboplatin OR as a single agent following multi-modality anthracycline-based therapy, B) HER2-overexpressing metastatic breast cancer AND medication will be used in combination with paclitaxel for first-line treatment OR as a single agent in a patient who received one or more chemotherapy regimens for metastatic disease OR in combination with Perjeta (pertuzumab) and docetaxel in a patient who has not received prior anti-HER2 therapy (e.g., trastuzumab) or chemotherapy for metastatic disease C) HER2 overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma AND patient has not received prior treatment for metastatic disease AND medication will be used in combination with cisplatin and capecitabine or 5-fluorouracil
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Subject to B vs D. Prescriber has assessed the patient's cardiac function/left ventricular ejection fraction prior to initiation of therapy. Female patients of child-bearing potential have been advised of the risk of fetal harm and the need for contraception.

HEXALEN

Products Affected

- Hexalen

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Severe bone marrow depression-indicated by CBC. Severe neurologic toxicity-Seizure.
Required Medical Information	Diagnosis of persistent or recurrent ovarian cancer AND the medication will be used as palliative treatment AND the medication will be used as a single agent AND the medication will be used following first-line therapy with a cisplatin and/or alkylating agent-based combination.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HRM-ANALGESICS

Products Affected

- Bupap Oral Tablet 50-300 MG
- Butalbital-Acetaminophen Oral Tablet 50-325 MG
- Butalbital-APAP-Caff-Cod Oral Capsule 50-325-40-30 MG

- Butalbital-APAP-Caffeine Oral Tablet 50-325-40 MG
- Butalbital-ASA-Caff-Codeine
- Meperidine HCl Injection Solution 100 MG/ML, 25 MG/ML
- Pentazocine-Naloxone HCl

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Analgesics: APAP/codeine,hydrocodone/APAP,hydrocodone/IBU,hydromorphone, methadone,morphine sulfate,oxycodone,oxycodone/APAP,oxycodone/ASA,oxycodone/ibuprofen,oxymorphone IR,tramadol, tramadol/APAP

HRM-ANTIARRHYTHMICS

Products Affected

- Digitek Oral Tablet 250 MCG
- Digox Oral Tablet 250 MCG
- Digoxin Injection
- Digoxin Oral Solution
- Digoxin Oral Tablet 250 MCG
- Disopyramide Phosphate Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Anti-arrhythmics: DIGOXIN: digoxin 0.125mg dose, propranolol, or sotalol for atrial fibrillation, DISOPYRAMIDE: dofetilide, amiodarone, propafenone, mexiletine, multaq

HRM-ANTICONVULSANTS

- Products Affected**
- PHENobarbital Oral Elixir
 - PHENobarbital Oral Tablet

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Anticonvulsants: Aptiom, Banzel, carbamazepine, Celontin, Cerebyx, clonazepam, diazepam, Dilantin, divalproex, ethosuximide, felbamate, fosphenytoin, Fycompa, gabapentin, gabitril, lamotrigine, levetiracetam, Lyrica, Onfi, oxcarbazine, Oxtellar, Peganone, phenytoin, Potiga, Primidone, Qudexy XR, Sabril, Tegretol-XR, Tiagabine, topiramate, Trokendi-XR, valproate, Vimpat, zonisamide

HRM-ANTIDEMENTIA

Products Affected

- Ergoloid Mesylates Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Antidementia: donepezil,galantamine,memantine,rivastigmine oral

HRM-ANTIDEPRESSANTS

Products Affected

- Amitriptyline HCl Oral
- Chlordiazepoxide-Amitriptyline
- ClomiPRAMINE HCl Oral
- Doxepin HCl Oral
- Imipramine HCl Oral
- Imipramine Pamoate
- Perphenazine-Amitriptyline
- Trimipramine Maleate Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Antidepressants: SSRI, SNRI, bupropion, mirtazapine, trazodone

HRM-ANTIHISTAMINES

Products Affected

- Carbinoxamine Maleate Oral Tablet 4 MG
- Cyproheptadine HCl Oral
- HydrOXYzine HCl Intramuscular
- HydrOXYzine HCl Oral Syrup
- HydrOXYzine HCl Oral Tablet
- HydrOXYzine Pamoate Oral
- Phenadoz Rectal Suppository 25 MG
- Promethazine HCl Injection
- Promethazine HCl Oral Syrup
- Promethazine HCl Oral Tablet
- Promethazine HCl Rectal Suppository 12.5 MG
- Promethegan Rectal Suppository 12.5 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None

HRM-ANTIINFLAMMATORY

Products Affected

- Indomethacin ER
- Indomethacin Oral

- Ketorolac Tromethamine Injection Solution 15 MG/ML, 30 MG/ML
- Ketorolac Tromethamine Intramuscular Solution 60 MG/2ML
- Ketorolac Tromethamine Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Anti-inflammatories: celecoxib, diclofenac, diflunisal, etodolac, flurbiprofen, ibuprofen, ketoprofen, meclofenamate, meloxicam, nabumetone, naproxen, oxaprozin, piroxicam, sulindac, tolmetin

HRM-ANTINEOPLASTICS

Products Affected

- Megestrol Acetate Oral Suspension 40 MG/ML, 625 MG/5ML
- Megestrol Acetate Oral Tablet

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	For Megestrol: dronabinol

HRM-ANTIPARKINSONS

Products Affected

- Bzotropine Mesylate Injection

- Bzotropine Mesylate Oral
- Trihexyphenidyl HCl

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Antiparkinsons': amantadine, Apokyn, Azilect, carbidopa/levodopa, entacapone, Neupro, p ramipexole, ropinirole, selegiline

HRM-ANTIPLATELET

Products Affected

- Dipyridamole Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Anti-platelets: Anagrelide, asa/dipyridamole, Brilinta, cilostazol, clopidogrel

HRM-ANTIPSYCHOTICS

- Products Affected**
- ChlorproMAZINE HCl Injection Solution 50 MG/2ML
 - ChlorproMAZINE HCl Oral
 - Perphenazine Oral
 - Thioridazine HCl Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Applies to New Starts only. Haloperidol, quetiapine, risperidone, aripiprazole, asenapine, iloperidone, lurasidone, olanzapine, paliperidone, ziprasidone

HRM-ANXIOLYTICS

Products Affected

- Meprobamate

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	SSRI, SNRI, buspirone.

HRM-CARDIOVASCULAR

Products Affected

- GuanFACINE HCl ER
- GuanFACINE HCl Oral
- Methyldopa Oral

- Methyldopa-Hydrochlorothiazide
- Methyldopate HCl
- NIFEdipine Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Cardiovascular agents: acebutolol, amilor/hctz, amlod/benazp, amlod/valsar, amlodipine, atenol/chlorth, atenolol, benazep/hctz, benazepril, benicar, benicar hct, betaxolol, bisopr/hctz, bisoprolol, candesartan, candesartan/hctz, captopril/hctz, captopril, cartia xt, carvedilol, chlorothiazide, diltiazem, dilt-xr, doxazosin, enalapril, enalapril/hctz, eprosartan, felodipine, fosinopril, fosinopril/hctz, hctz, indapamide, irbesart/hctz, irbesartan, isradipine, labetalol, lisinopril, lisinopril/hctz, losartan/losartan/hctz, methylclothia, metolazone, meto

PA Criteria	Criteria Details
	prol/hctz, metoprolol, midodrine, moexipril/hctz, moexipril, nadolol, nadolol/bend, nifedipine, nifedical xl, nimodipine, nifedipine er, nisoldipine, perindopril, pindolol, prazosin, propranolol, quinapril/quinapril/hctz, ramipril, spirono/hctz, taztia xt, telmis/amlod, telmis/hctz, telmisartan, terazosin, timolol, trandolapril, trandolapril/verapamil, triam/hctz, valsart/hctz, valsartan, verapamil

HRM-ORAL AND TRANSDERMAL ESTROGENS AND PROGESTINS

Products Affected

- Divigel Transdermal Gel 1 MG/GM
- Elestrin
- Estradiol Oral
- Estradiol Transdermal
- Estropipate Oral Tablet 0.75 MG
- Evamist
- Menest Oral Tablet 0.3 MG, 0.625 MG, 1.25 MG
- Norethindrone-Eth Estradiol

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Bone Density: alendronate, risedronate, ibandronate, raloxifene (zoledronic acid for bed-bound patients or for post-hip fracture). Vaginal Symptoms: vaginal estrogen cream

HRM-SEDATIVE HYPNOTICS

Products Affected

- Butisol Sodium Oral Tablet 30 MG
- Zaleplon Oral Capsule 10 MG, 5 MG
- Zolpidem Tartrate ER
- Zolpidem Tartrate Oral Tablet 10 MG, 5 MG
- Zolpidem Tartrate Sublingual

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Sleep disorder agents: estazolam, flurazepam, rozerem ,temazepam, triazolam

HRM-SKELETAL MUSCLE RELAXANTS

Products Affected

- Carisoprodol Oral Tablet 350 MG
- Chlorzoxazone Oral Tablet 500 MG
- Cyclobenzaprine HCl Oral Tablet 10 MG, 5 MG
- Methocarbamol Injection Solution 1000 MG/10ML
- Methocarbamol Oral
- Orphenadrine Citrate ER
- Orphenadrine Citrate Injection

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HRM-SULFONYLUREAS

Products Affected

- ChlorproPAMIDE
- GlyBURIDE Micronized
- GlyBURIDE Oral
- GlyBURIDE-MetFORMIN

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	High risk medication. Automatically approved for beneficiaries less than or equal to 64 years. Attestation that risk outweighs benefit as to the medical necessity for using this high risk medication and anticipated treatment course/duration, and if formulary alternatives considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 alternative(s) - See OTHER criteria for alternatives.
Age Restrictions	Less than or equal to 64 years old, claim for target drug automatically pays. Greater than or equal to 65 years old, prior authorization exception request is required indicating medically accepted indication not otherwise excluded from Part D.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Anti-Diabetics: glimepiride, glipizide IR and ER

HUMIRA

Products Affected

- Humira Pediatric Crohns Start Subcutaneous Prefilled Syringe Kit
- Humira Pen Subcutaneous Pen-Injector Kit
- Humira Pen-CD/UC/HS Starter
- Humira Pen-Ps/UV Starter
- Humira Subcutaneous Prefilled Syringe

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Active serious infection (including tuberculosis). Combined use with a biologic disease-modifying anti-rheumatic drugs or potent immunosuppressant (e.g., azathioprine or cyclosporine)
Required Medical Information	Diagnosis of ONE of the following: A) moderate to severe rheumatoid arthritis OR moderate to severe polyarticular juvenile idiopathic arthritis and patient had inadequate response, intolerance, or contraindication to one or more non-biologic DMARDs for at least 3 consecutive months B) psoriatic arthritis and patient had inadequate response, intolerance, or contraindication to methotrexate C) ankylosing spondylitis and patient had inadequate response, intolerance, or contraindication to one or more NSAIDs D) moderate to severe chronic plaque psoriasis (affecting more than 5% of body surface area or affecting crucial body areas such as the hands, feet, face, or genitals) and patient had inadequate response, intolerance, or contraindication to conventional therapy with at least one of the following: phototherapy (including but not limited to UVA with a psoralen [PUVA] and/or retinoids [RePUVA] for at least one continuous month or one or more oral systemic treatments (Cyclosporine, acitretin, sulfasalazine, methotrexate, leflunomide, azathioprine) for at least 3 consecutive months E) moderate to severe Crohn's disease and patient had inadequate response, intolerance, or contraindication to conventional therapy with two or more of the following: corticosteroids or non-biologic DMARDs F) moderate to severe ulcerative colitis and patient had inadequate response, intolerance or contraindication to conventional therapy with two or more of the following: corticosteroids, 5-ASA (i.e. mesalamine, sulfasalazine, balsalazide) or non-biologic DMARDs (azathioprine, cyclosporine, hydroxychloroquine, leflunomide, penicillamine, sulfasalazine) G) non-infectious uveitis (including intermediate, posterior, and panuveitis) and patient had inadequate

PA Criteria	Criteria Details
	response, intolerance or contraindication to conventional therapy with one of the following following: systemic or topical corticosteroids or ophthalmic antimuscarinics. OR H) moderate to severe hidradenitis suppurativa
Age Restrictions	2 years of age or older for JIA. 6 years of age or older for pediatric Crohn's disease, 18 years of age or older for all other indications
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

HYDROXYPROGESTERONE

Products Affected

- Hydroxyprogesterone Caproate Intramuscular Solution

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Breast, cervical, hepatocellular, uterine, or vaginal cancers, hepatic or thromboembolic disease, jaundice, or vaginal bleeding
Required Medical Information	Supporting statement of diagnosis from physician
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	21 weeks
Other Criteria	None

IBRANCE

Products Affected

- Ibrance

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer in combination with an aromatase inhibitor in postmenopausal women as initial endocrine-based therapy OR in combination with fulvestrant in women with disease progression following endocrine therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	None

ICLUSIG

Products Affected

- Iclusig

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic myelogenous leukemia(CML) AND One of the following: A) History of failure, resistance, relapse, contraindication, or intolerance to at least TWO other tyrosine kinase inhibitors (i.e., GLEEVEC [imatinib], SPRYCEL, TASIGNA, and BOSULIF), or B) Patient has the T315I mutation. OR Diagnosis of Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) AND One of the following: A) History of failure, resistance relapse, contraindication, or intolerance to at least TWO other FDA-approved tyrosine kinase inhibitors (i.e., GLEEVEC [imatinib], SPRYCEL), or B) Patient has the T315I mutation.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None

IDHIFA

Products Affected

- IDHIFA

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia with an isocitrate dehydrogenase 2 mutation as detected by an FDA approved test.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of therapy

IMATINIB

Products Affected

- Imatinib Mesylate

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML), B) Ph+ acute lymphoblastic leukemia (ALL), C) Gastrointestinal tumor (GIST) where patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST or patient had resection of c-KIT positive GIST and imatinib will be used as an adjuvant therapy, D) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic, E) hypereosinophilic syndrome or chronic eosinophilic leukemia, F) myelodysplastic syndrome or myeloproliferative disease associated with platelet-derived growth factor receptor gene rearrangements, G) aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutational status unknown
Age Restrictions	1 year of age or older - newly diagnosed CML in the chronic phase or newly diagnosed Ph+ ALL. 18 years of age or older for other indications.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

IMBRUVICA

Products Affected

- Imbruvica

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of mantle cell lymphoma (MCL) who have received at least one prior therapy OR chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) OR chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) with 17p deletion OR Waldenstrom's macroglobulinemia (WM) OR marginal zone lymphoma who require systemic therapy and have received at least one prior anti-CD20-based therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

IMFINZI

Products Affected

- Imfinzi

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic urothelial carcinoma. Patient must have progressed on or following platinum-containing chemotherapy, or within 12 months of neoadjuvant or adjuvant platinum containing chemotherapy OR Unresectable Stage III, non-small cell lung cancer without progression following concurrent platinum-based chemotherapy and radiation therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a oncologist
Coverage Duration	12 months
Other Criteria	None

INCRELEX

Products Affected

- Increlex

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Increlex is contraindicated in patients with allergies to mecasermin or any component of the Increlex formulation, for growth promotion in patients with closed epiphyses, for IV administration, in patients with active or suspected neoplasia. Increlex should be discontinued if neoplasia develops while on therapy.
Required Medical Information	Increlex (mecasermin [rDNA origin] injection) is indicated for the long-term treatment of growth failure in children with severe primary IGF-1 deficiency (Primary IGFD) or with growth hormone (GH) gene deletion who have developed neutralizing antibodies to GH. Child has one of the following conditions: Severe primary IGF-1 deficiency, OR Growth hormone gene deletion with developed neutralizing antibodies to growth hormone, OR Genetic mutation of GH receptor (i.e. Laron Syndrome), AND Child has severe growth retardation with height standard deviation score (SDS) more than 3 SDS below the mean for chronological age and sex, AND Child with IGF-1 level greater than or equal to 3 standard deviations below normal based on lab reference range for age and sex, AND Child with normal or elevated growth hormone (GH) levels based on at least one growth hormone stimulation test, AND Evidence of open epiphyses
Age Restrictions	2 years of age and older
Prescriber Restrictions	Pediatric or Endocrinologist
Coverage Duration	6 months
Other Criteria	For renewal, patient has experienced improvement

INHALED TOBRAMYCIN

Products Affected

- Tobi Podhaler

- Tobramycin Inhalation

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis AND patient has evidence of P. aeruginosa in the lungs
Age Restrictions	6 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations).

INLYTA

Products Affected

- Inlyta

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced renal cell carcinoma AND patient failed one or more systemic therapies for renal cell carcinoma (e.g., sunitinib-, bevacizumab-, temsirolimus-, or cytokine-containing regimens)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

INTRAROSA

Products Affected

- Intrarosa

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Vaginal bleeding or dysfunctional uterine bleeding of an undetermined origin, known or suspected estrogen-dependent neoplasia.
Required Medical Information	Diagnosis of moderate to severe dyspareunia or atrophic vaginitis AND A) Patient must be female, B) Patient must be menopausal or postmenopausal, C) Patient has tried and failed, has a contraindication or intolerance to a low dose vaginal estrogen preparation (e.g. Premarin vaginal cream, Estrace vaginal cream, Estring, Vagifem), D) Patient does not have renal or hepatic impairment.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 3 months, Reauthorization: 12 months
Other Criteria	None

INTRON A

Products Affected

- Intron A

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Uncontrolled depression. Solid organ transplant other than liver. Autoimmune hepatitis or other autoimmune condition known to be exacerbated by interferon and ribavirin.
Required Medical Information	Diagnosis of hairy cell leukemia OR Diagnosis of Condylomata acuminata OR Diagnosis of AIDS-related Kaposi's sarcoma OR Clinically aggressive follicular lymphoma and the medication will be used concurrently with anthracycline-containing chemotherapy or is not a candidate for anthracycline-containing chemotherapy OR Malignant melanoma and the request for coverage is within 56 days of surgery and the patient is at high risk of disease recurrence OR Diagnosis of chronic hepatitis B with compensated liver disease and patient has evidence of hepatitis B viral replication and patient has been serum hepatitis B surface antigen-positive for at least 6 months OR Diagnosis of chronic hepatitis C with compensated liver disease and is receiving combination therapy with ribavirin, unless ribavirin is contraindicated, and the medication will not be used as part of triple therapy with a protease inhibitor and patient has a clinical reason for not using peginterferon
Age Restrictions	1 year of age or older for HBV. 3 years of age or older for HCV. 18 years of age or older for other indications.
Prescriber Restrictions	None
Coverage Duration	Condylomata: 3 mos. HBV e antigen pos: 16 wks, e antigen neg: 48 wks. KS: 16 wks. Others: 12 mos
Other Criteria	None

IRESSA

Products Affected

- Iressa

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC AND Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ISTODAX

Products Affected

- Istodax (Overfill)

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma (CTCL) in patients who have received at least 1 prior systemic therapy, i.e. retinoid (acitretin, bexarotene) OR peripheral T-cell lymphoma (PTCL) in patients who have received at least 1 prior therapy.
Age Restrictions	None
Prescriber Restrictions	none
Coverage Duration	12 months
Other Criteria	None

JAKAFI

Products Affected

- Jakafi

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Myelofibrosis: Diagnosis of primary myelofibrosis, OR post-polycythemia vera myelofibrosis, OR post-essential thrombocythemia myelofibrosis. Polycythemia vera: Diagnosis of polycythemia vera, AND history of failure, contraindication, or intolerance to hydroxyurea.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None

JUXTAPID

Products Affected

- Juxtapid

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Moderate to severe liver impairment or active liver disease including unexplained persistent abnormal liver function tests. Pregnancy. Concomitant use with strong or moderate CYP3A4 inhibitors.
Required Medical Information	Diagnosis of HoFH as confirmed by one of the following: a) genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (ie, LDLRAP1 or ARH), or b) both of the following: 1) either untreated LDL-C greater than 500 mg/dL or treated LDL-C greater than 300 mg/dL AND 2) either xanthoma before 10 years of age or evidence of heterozygous FH in both parents. Patient has been receiving at least 12 consecutive weeks of ONE LDL-C lowering prescription therapy and will continue to receive an LDL-C lowering prescription therapy. History of failure after 12 consecutive weeks or intolerance to PCSK9 inhibitor therapy. One of the following LDL-C values while on maximally tolerated lipid-lowering regimen within the last 30 days: LDL-C greater than or equal to 100 mg/dL with ASCVD OR LDL-C greater than or equal to 130 mg/dL without ASCVD. Not used in combination with Kynamro (mipomersen). Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor. Patient is not pregnant. Patient does not have moderate or severe hepatic impairment (ie, Child-Pugh category B or C) or active liver disease including unexplained persistent abnormal liver function tests. Patient is not concomitantly on moderate or strong CYP 3A4 inhibitors (eg, clarithromycin).
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	HoFH (initial): 6 months. (reauth): 12 months
Other Criteria	None

KADCYLA

Products Affected

- Kadcyła

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Patient has a diagnosis of HER2-positive metastatic breast cancer and the member has been previously treated with trastuzumab and a taxane
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Prescriber has assessed the patient's hepatic function and left ventricular ejection fraction prior to initiation of therapy. Female patients of child-bearing potential had pregnancy status verified prior to the initiation of Kadcyła and have been advised of the risk of fetal harm and the need for contraception.

KALYDECO

Products Affected

- Kalydeco

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis AND the patient has 1 mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data.
Age Restrictions	For oral granules- 2 years of age or older. For oral tablets- 6 years of age or older.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient has experienced benefit from therapy (i.e. improvement in pulmonary lung function [FEV1], decreased number of pulmonary exacerbations)

KANUMA

Products Affected

- Kanuma

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Lysosomal acid lipase deficiency
Age Restrictions	None
Prescriber Restrictions	Prescribed by hepatologist
Coverage Duration	12 months
Other Criteria	None

KEYTRUDA

Products Affected

- Keytruda Intravenous Solution

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	<p>Diagnosis of unresectable or metastatic melanoma OR first-line treatment of metastatic non-small cell lung cancer (NSCLC) in patients with high PD-L1 expressing tumors and with no EGFR or ALK genomic tumor aberrations OR treatment of metastatic NSCLC in patients with PD-L1 expression who have disease progression on or after platinum-containing chemotherapy (patients with EGFR or ALK genomic tumor aberrations should have disease progression on approved therapy for these aberrations prior to receiving Keytruda) OR recurrent or metastatic cervical cancer with tumor PD-L1 expression and disease progression on or after previous chemotherapy OR recurrent or metastatic squamous cell carcinoma of the head and neck in patients with disease progression on or after platinum-containing chemotherapy OR treatment of adult or pediatric patients with classical Hodgkin lymphoma (in patients who are refractory or who have relapsed after 3 or more prior lines of therapy) OR first-line treatment (in combination with pemetrexed plus carboplatin) of metastatic nonsquamous NSCLC OR locally advanced or metastatic urothelial carcinoma (in patients who are not eligible for cisplatin-containing chemotherapy, or who have had disease progression during or following platinum-containing chemotherapy or within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy) OR unresectable or metastatic solid tumors that have been identified as having a biomarker referred to as microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) OR for the treatment of patients with recurrent locally advanced or metastatic gastric or gastroesophageal junction adenocarcinoma whose tumors express PD-L1 as determined by an FDA-approved test, with disease progression on or after two or more prior lines of therapy including fluoropyrimidine and platinum-containing chemotherapy and if appropriate, HER2/neu-targeted therapy.</p>

PA Criteria	Criteria Details
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

KINERET

Products Affected

- Kineret Subcutaneous Solution Prefilled Syringe

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Active infection, concurrent therapy with other biologics.
Required Medical Information	Must have one of following: 1) inadequate response to methotrexate (MTX), 2) inadequate response to another nonbiologic DMARD (e.g., leflunomide, hydroxychloroquine, sulfasalazine) if contraindicated or intolerant to MTX, 3) intolerance or contraindication to at least 2 nonbiologic DMARDs or, 4) use Kineret as first-line therapy with MTX for severely active RA. For diagnosis of CAPs, Kineret will be approved. For rheumatoid arthritis member needs trial or intolerance/contraindication to Humira and Enbrel.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For re-authorization, patient's condition must have improved or stabilized.

KISQALI

Products Affected

- Kisqali 200 Dose
- Kisqali 400 Dose
- Kisqali 600 Dose
- Kisqali Femara 200 Dose
- Kisqali Femara 400 Dose
- Kisqali Femara 600 Dose

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR)-positive, HER2-negative advanced or metastatic breast cancer and intended to be used in combination with an aromatase inhibitor in postmenopausal women.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a oncologist
Coverage Duration	12 months
Other Criteria	None

KORLYM

Products Affected

- Korlym

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy
Required Medical Information	Supporting statement of diagnosis and relevant medical information from physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

KUVAN

Products Affected

- Kuvan

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of hyperphenylalaninemia (HPA) due to tetrahydrobiopterin (BH4)-responsive phenylketonuria (PKU). Appropriate for use in patients who: a) have been diagnosed with PKU, b) have a baseline blood Phe measured within 2 weeks prior to initiating therapy. Also require that the prescriber be a specialist with knowledge and expertise in metabolic diseases or genetic diseases or has consulted with a specialist in metabolic or genetic diseases. Initial approval will be for two months of therapy if the initial dose is 5 mg/kg/day to less than 20 mg/kg/day, it will be for one month if the initial dose is 20 mg/kg/day. Renewal for continued use will be for 6 months if patient response is seen based on prescriber determination.
Age Restrictions	1 month of age or older
Prescriber Restrictions	Specialist knowledgeable in the management of PKU
Coverage Duration	Initial Approval: 2 months. Extended Approval: 6 month intervals
Other Criteria	Blood Phe levels should be checked after 1 week of therapy and periodically up to one month during a therapeutic trial.

KYNAMRO

Products Affected

- Kynamro Subcutaneous Solution Prefilled Syringe

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Moderate to severe liver impairment or active liver disease including unexplained persistent abnormal liver function tests.
Required Medical Information	Diagnosis of homozygous familial hypercholesterolemia as evidenced by one of the following: A) genetic confirmation of 2 mutant alleles at the LDL receptor, ApoB, PCSK9, or autosomal recessive hypercholesterolemia (ARH) adaptor protein gene locus OR B) untreated/pre-treatment LDL greater than 500 mg/dL with at least one of the following: cutaneous or tendonous xanthoma before age 10 years, history of early vascular disease (men younger than 55 years, women younger than 60 years) on both sides of the family if parenteral LDL levels are unknown, elevated LDL cholesterol levels before lipid-lowering therapy consistent with heterozygous FH in both parents AND Patient has tried and had an inadequate response to the maximum tolerated dose of a high potency statin (e.g., atorvastatin, rosuvastatin), unless all statins are contraindicated.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial - 6 months. Renewal - 12 months.
Other Criteria	For renewal, patient has responded to therapy with a decrease in LDL levels from baseline AND patient does not have contraindications to therapy.

LARTRUVO

Products Affected

- Lartruvo

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of soft tissue sarcoma (STS), histologic subtype for which an anthracycline-containing regimen is appropriate, previous treatment failure with radiotherapy or surgery and must document being used in combination with doxorubicin for the first 8 cycles.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

LENVIMA

Products Affected

- Lenvima 10 MG Daily Dose
- Lenvima 14 MG Daily Dose
- Lenvima 18 MG Daily Dose
- Lenvima 20 MG Daily Dose
- Lenvima 24 MG Daily Dose
- Lenvima 8 MG Daily Dose

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally recurrent or metastatic, progressive, radioactive iodine-refractory differentiated thyroid cancer OR advanced renal cell carcinoma following one prior anti-angiogenic therapy in combination with everolimus
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

LETAIRIS

Products Affected

- Letairis

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND Patient has WHO Group I PAH AND pregnancy must be excluded prior to the start of therapy and will be prevented thereafter with two forms of reliable contraception in female patients of reproductive potential.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial - 6 months. Renewal - 12 months
Other Criteria	None

LEUKINE

Products Affected

- Leukine Intravenous

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Concurrent use with myelosuppressive chemotherapy or radiation or excessive (greater than or equal to 10%) leukemic myeloid blasts in bone marrow or peripheral blood
Required Medical Information	Diagnosis of one of the following: A) Patient has undergone allogeneic or autologous bone marrow transplant (BMT) and engraftment is delayed or failed OR B) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis OR C) Medication will be used for myeloid reconstitution after an autologous or allogeneic BMT OR D) Patient has acute myeloid leukemia and administration will be after completion of induction chemotherapy
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

LEUPROLIDE

Products Affected

- Eligard
- Leuprolide Acetate Injection
- Lupron Depot (1-Month)
- Lupron Depot (3-Month)
- Lupron Depot (4-Month)
- Lupron Depot (6-Month)
- Lupron Depot-Ped (1-Month) Intramuscular Kit 11.25 MG, 15 MG
- Lupron Depot-Ped (3-Month) Intramuscular Kit 30 MG (Ped)

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) advanced or metastatic prostate cancer, B) Diagnosis of central precocious puberty and patient had early onset of secondary sexual characteristics (male: earlier than 9 years of age. female: earlier than 8 years of age) and advanced bone age of at least one year compared with chronological age and has undergone gonadotropin-releasing hormone agonist (GnRHa) testing with peak luteinizing hormone (LH) level above pre-pubertal range or random LH level in pubertal range and Patient had the following diagnostic evaluations to rule out tumors, when suspected: diagnostic imaging of the brain (MRI or CT scan), Pelvic/testicular/adrenal ultrasound, Human chorionic gonadotropin levels, Adrenal steroids to rule out congenital adrenal hyperplasia, C) the medication will be used for stimulation testing to confirm the diagnosis of central precocious puberty or D) management of endometriosis
Age Restrictions	None
Prescriber Restrictions	CPP - Prescribed by or in consultation with a pediatric endocrinologist
Coverage Duration	12 months. CPP testing: one time dose.
Other Criteria	For renewal of CPP, LH levels have been suppressed to pre-pubertal levels and consideration for discontinuation of therapy when the patient is 11 years of age for girls and 12 years of age for boys.

LIDOCAINE PATCH

Products Affected

- Lidocaine External Patch 5 %

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of pain associated with diabetic neuropathy OR pain associated with cancer-related neuropathy OR post-herpetic neuralgia.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

LINEZOLID

Products Affected

- Linezolid Intravenous Solution 600 MG/300ML
- Linezolid Oral Tablet

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Not covered with concomitant use of MAOI therapy
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	VRE: 4 weeks. Nosocomial and community acquired pneumonia: 3 weeks. All other indications: 2 weeks
Other Criteria	None

LONSURF

Products Affected

- Lonsurf

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic colorectal cancer, previously treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based regimens, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For initial treatment: Absolute neutrophil count 1,500/mm ³ or greater or febrile neutropenia resolved, platelet count 75,000/mm ³ or greater, and grade 3 or 4 nonhematological reactions resolved to grade 0 or 1

LYNPARZA

Products Affected

- Lynparza

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Appropriate diagnosis and testing for BRCA mutation (deleterious or suspected deleterious germline BRCA mutated (as detected by an FDA approved test) advanced ovarian cancer that has been treated with 3 or more prior lines of chemotherapy)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

MEKINIST

Products Affected

- Mekinist

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic melanoma and medication is used as a single agent and patient has a positive BRAF V600E or V600K mutation as detected by an FDA-approved test (THxID-BRAF Kit) or Clinical Laboratory Improvement Amendments (CLIA)-approved facility, and the patient has not received prior BRAF-inhibitor therapy (i.e. Zelboraf, Tafinlar) OR medication will be used in combination with Tafinlar in a patient with BRAF V600E or V600K mutations, as detected by an FDA-approved test (THxID-BRAF kit) or Clinical Laboratory Improvement Amendments (CLIA)-approved facility.
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

MOZOBIL

Products Affected

- Mozobil

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Patient is to undergo autologous stem cell transplantation for the treatment of non-Hodgkin's lymphoma or multiple myeloma AND Patient will concomitantly receive a daily dose of a granulocyte colony-stimulating factor (G-CSF) for 4 days prior to the first evening dose of Mozobil and on each day prior to apheresis while using Mozobil.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	4 days
Other Criteria	None

MS INTERFERONS

Products Affected

- Betaseron Subcutaneous Kit

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsing form of multiple sclerosis OR diagnosis of first clinical episode and MRI features consistent with multiple sclerosis.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient has experienced an objective response to therapy (i.e. no or slowed progression of disease)

MYLOTARG

Products Affected

- Mylotarg Intravenous Solution Reconstituted 4.5 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	INITIAL: A. Newly- diagnosed, CD33 positive acute myeloid leukemia (AML) or B. Relapsed or refractory CD33 positive AML. CONTINUATION OF THERAPY: 1) patient continues to meet initial criteria and 2) patients with newly diagnosed AML have not exceeded a maximum of 8 cycles
Age Restrictions	Relapsed or refractory AML: 2 years and older, Newly diagnosed AML: 18 years and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None

MYTESI

Products Affected

- Mytesi

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	CLINICAL NOTES TO SUPPORT A DIAGNOSIS OF CHRONIC DIARRHEA, DEFINED AS DIARRHEA PERSISTING FOR MORE THAN FOUR WEEKS, CAUSED BY THEIR MEDICATION REGIMEN OR HIV ENTEROPATHY PROVEN BY BIOPSY, AND NOT A VIRUS, PARASITE OR BACTERIUM AS EVIDENCED BY STOOL SAMPLE TAKEN IN THE PREVIOUS 3 MONTHS. PATIENT MUST HAVE TRIED AND FAILED OR HAD INTOLERANCE TO LOPERAMIDE OR DIPHENOXYLATE-ATROPINE TRIALS OF A MINIMUM OF 30 DAYS.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Infectious Disease Specialist or GI Consult for new starts
Coverage Duration	12 months
Other Criteria	None

NAGLAZYME

Products Affected

- Naglazyme

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Mucopolysaccharidosis VI (MPS VI or Maroteaux-Lamy syndrome).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NATPARA

Products Affected

- Natpara

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Statement of diagnosis from the prescriber
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NERLYNX

Products Affected

- Nerlynx

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of early stage HER2- overexpressed breast cancer. Must be used after trastuzumab therapy.
Age Restrictions	age 18 years and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None

NEUPOGEN

Products Affected

- Neupogen Injection Solution 300 MCG/ML, 480 MCG/1.6ML

- Neupogen Injection Solution Prefilled Syringe
- Zarxio

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	<p>Diagnosis of one of the following: A) congenital, cyclic, or idiopathic neutropenia, B) severe febrile neutropenia (FN) with the following: Has not received prophylactic pegfilgrastim and Used as adjunct to appropriate antibiotics in high-risk patients and any one of the following: 65 years or older, Uncontrolled primary disease, Pneumonia, Hypotension and multiorgan dysfunction (sepsis syndrome), Invasive fungal infection, Hospitalization when developed fever, Prior FN, Severe (ANC less than 100/mcL) or anticipated prolonged (more than 10 days) neutropenia, C) Autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, D) Undergoing myeloablative chemotherapy followed by autologous or allogeneic BMT, E) Acute myeloid leukemia and will be given after completion of induction or consolidation chemotherapy, F) Acute lymphoblastic leukemia and will be given after completion of the first few days of chemotherapy of the initial induction or first post-remission course, G) Myelodysplastic syndrome with severe neutropenia and recurrent infection, H) Receiving radiation therapy, not on chemotherapy, and expected to have prolonged delays in treatment due to neutropenia, I) Neutropenia associated with HIV infection and antiretroviral therapy, J) Aplastic anemia, K) Primary prophylaxis of FN in one of the following patients: 20% or higher risk of FN based on chemotherapy regimen OR Less than 20% risk of FN based on chemotherapy regimen with one of the following: 65 years or older, Poor performance status, Poor nutritional status, Previous FN, Extensive prior treatment including large radiation ports, Cytopenias due to bone marrow involvement by tumor, Administration of combined chemoradiotherapy, Presence of open wounds or active infections, Other serious comorbidities (including renal or liver dysfunction) or Receiving dose-dense chemotherapy regimen in</p>

PA Criteria	Criteria Details
	breast or small cell lung cancer or non-Hodgkins lymphoma.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NEXAVAR

Products Affected

- NexAVAR

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Squamous cell lung cancer being treated with carboplatin and paclitaxel.
Required Medical Information	Diagnosis of unresectable hepatocellular carcinoma OR Diagnosis of advanced renal cell carcinoma OR Diagnosis of locally recurrent or metastatic, progressive, differentiated thyroid carcinoma refractory to radioactive iodine treatment
Age Restrictions	18 years and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NINLARO

Products Affected

- Ninlaro

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple myeloma, documentation of combination therapy with lenalidomide and dexamethasone. History of 1 prior therapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NORTHERA

Products Affected

- Northera

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Prior authorization will be approved for the following indication(s): orthostatic dizziness, light-headedness, or the feeling that you are about to black out in adults with neurogenic orthostatic hypotension (NOH) caused by primary autonomic failure (i.e., Parkinson disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, and non-diabetic autonomic neuropathy)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NOXAFIL

Products Affected

- Noxafil Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Concomitant treatment with sirolimus, CYP 3A4 substrates that prolong QT interval (pimozide, quinidine), HMG-CoA Reductase inhibitors primarily metabolized through CYP 3A4, or ergot alkaloids
Required Medical Information	Diagnosis of oropharyngeal candidiasis and patient tried itraconazole and/or fluconazole OR Medication will be used as prophylaxis of invasive Aspergillus and Candida infections and the patient is at high risk of developing these infections due to being severely immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with graft-versus-host disease (GVHD) or those with hematologic malignancies with prolonged neutropenia from chemotherapy.
Age Restrictions	13 years of age or older for prophylaxis of invasive aspergillus or candidal infection
Prescriber Restrictions	None
Coverage Duration	12 weeks
Other Criteria	None

NUCALA

Products Affected

- Nucala

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of severe asthma (eosinophilic phenotype) OR eosinophilic granulomatosis with polyangiitis (EGPA)
Age Restrictions	12 years of age or older for severe asthma eosinophilic phenotype or 18 years of age or older for granulomatosis with polyangiitis (EGPA)
Prescriber Restrictions	Must be prescribed by a pulmonologist or immunologist
Coverage Duration	12 months
Other Criteria	None

NUEDEXTA

Products Affected

- Nuedexta

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of pseudobulbar affect.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

NULOJIX

Products Affected

- Nulojix

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Kidney transplant: The medication is being used for prevention of kidney transplant organ rejection AND The patient is immune to the Epstein-Barr virus (i.e. EBV seropositive) AND The patient is prescribed concurrent therapy with mycophenolate and corticosteroids
Age Restrictions	Kidney transplant: 18 years of age or older
Prescriber Restrictions	Kidney transplant: Prescriber is experienced in immunosuppressive therapy and management of transplant patients
Coverage Duration	12 months
Other Criteria	Subject to Part B vs. Part D review. Approve for continuation of prior therapy.

NUPLAZID

Products Affected

- Nuplazid Oral Tablet 17 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Parkinson disease psychosis including hallucinations and/or delusions
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ODOMZO

Products Affected

- Odomzo

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced basal cell carcinoma of the skin and specific documentation of negative pregnancy status
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

OPDIVO

Products Affected

- Opdivo Intravenous Solution 100 MG/10ML, 40 MG/4ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of BRAF V600 wild-type or BRAF V600 mutation-positive unresectable or metastatic melanoma and used as single agent OR unresectable or metastatic melanoma in combination with ipilimumab OR for the adjuvant treatment of melanoma in patients with lymph node involvement or metastatic disease who have undergone complete resection OR treatment of patients with metastatic non-small cell lung cancer (NSCLC) with progression on or after platinum-based chemotherapy and patients with EGFR or ALK genomic tumor aberrations should have disease progression (on FDA-approved EGFR- or ALK-directed therapy) prior to receiving nivolumab OR advanced renal cell carcinoma in combination with Yervoy OR as monotherapy in patients who have received prior anti-angiogenic therapy OR recurrent or metastatic squamous cell carcinoma of the head and neck in patients with disease progression on or after platinum-based chemotherapy OR classical Hodgkin lymphoma in patients who have relapsed or progressed after 3 or more lines of systemic therapy that includes an autologous hematopoietic stem cell transplant (HSCT) and brentuximab vedotin OR locally advanced or metastatic urothelial carcinoma in patients with disease progression on or following platinum-containing therapy or within 12 months of neoadjuvant or adjuvant treatment with a platinum-containing therapy OR microsatellite instability-high or mismatch repair deficient metastatic colorectal cancer with progression after treatment of fluoropyrimidine, oxaliplatin, and irinotecan OR treatment of hepatocellular cancer, after disease progression on or intolerance to sorafenib therapy.
Age Restrictions	None
Prescriber Restrictions	None

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	None

OPSUMIT

Products Affected

- Opsumit

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension WHO group I AND diagnosis was confirmed by right heart catheterization AND female patients are enrolled in the OPSUMIT REMS program.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months - initial. 12 months - renewal
Other Criteria	None

ORENCIA

Products Affected

- Orenzia ClickJect
- Orenzia Intravenous

- Orenzia Subcutaneous Solution Prefilled Syringe

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Active infection (including TB). Concurrent therapy with other biologics.
Required Medical Information	Screening for latent TB infection and assessment for Hep B risk. For positive latent TB, patient must have completed treatment or is currently receiving treatment for LTBI. HBV infection ruled out or treatment initiated for positive infection. Rheumatoid arthritis - Must have one of following: 1) inadequate response to methotrexate (MTX), 2) inadequate response to another nonbiologic DMARD (e.g., leflunomide, hydroxychloroquine, sulfasalazine) if contraindicated or intolerant to MTX, 3) intolerance or contraindication to at least 2 nonbiologic DMARDs or, 4) use Orenzia as first-line therapy with MTX for severely active RA. Polyarticular JIA - Must have an inadequate response to at least one nonbiologic DMARD or intolerance/contraindication to at least 2 nonbiologic DMARDs. For PAJIA, member needs trial or intolerance/contraindication to Humira. For RA, member needs trial or intolerance/contraindication to Humira and Enbrel.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For re-authorization, patient's condition must have improved or stabilized.

ORKAMBI

Products Affected

- Orkambi Oral Tablet

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Initial Therapy: Diagnosis of cystic fibrosis (CF) with documented homozygous F508del mutation confirmed by FDA-approved CF mutation test AND if less than 18 years of age, baseline ophthalmological exam completed. Continuation of therapy: Documentation patient is tolerating and responding to medication (i.e. improved FEV1, weight gain, decreased exacerbations, etc.)
Age Restrictions	Must be greater than or equal to 12 years of age
Prescriber Restrictions	Must be prescribed by, or in conjunction with, a pulmonologist or is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None

OSPHEANA

Products Affected

- Osphena

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Vaginal bleeding or dysfunctional uterine bleeding of an undetermined origin, known or suspected estrogen-dependent neoplasia, acute thromboembolism or a past history of thromboembolic disease (including patients with a history of DVT, pulmonary embolism, retinal vein thrombosis, stroke, or myocardial infarction, known or suspected pregnancy.
Required Medical Information	Diagnosis of moderate to severe dyspareunia or atrophic vaginitis AND A) Patient must be female, B) Patient must be menopausal or postmenopausal, C) Patient has tried and failed, has a contraindication or intolerance to a low dose vaginal estrogen preparation (e.g. Premarin vaginal cream, Estrace vaginal cream, Estring, Vagifem), D) Dose must not exceed 1 tablet per day, E) Patient does not have hepatic impairment.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 2 months, Reauthorization: 12 months
Other Criteria	None

OTREXUP

Products Affected

- Otrexup Subcutaneous Solution Auto-Injector 10 MG/0.4ML, 12.5 MG/0.4ML, 15 MG/0.4ML, 20 MG/0.4ML, 22.5 MG/0.4ML, 25 MG/0.4ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy, breastfeeding, alcoholism, alcoholic liver disease or other chronic liver disease, immunodeficiency syndrome, or preexisting blood dyscrasias
Required Medical Information	Diagnosis of Psoriasis or Rheumatoid Arthritis, including polyarticular juvenile idiopathic arthritis, AND documented trial and failure, contraindication, or intolerance to oral methotrexate.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

OXANDROLONE

Products Affected

- Oxandrolone Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Breast or prostate cancer in men. Breast cancer in women with hypercalcemia. Pregnancy. Nephrosis or nephrotic phase of nephritis. Hypercalcemia.
Required Medical Information	Patient is receiving treatment as an adjunct therapy to promote weight gain and has one of the following: Extensive surgery, Chronic infections, Severe trauma, Failure to gain or maintain at least 90% of ideal body weight without definite pathophysiologic reasons and Patient has had an inadequate response, intolerance, or contraindication to nutritional supplements and a nutritional consult was performed OR Oxandrin (oxandrolone) will be used to counterbalance protein catabolism associated with chronic corticosteroid administration OR Patient has bone pain associated with osteoporosis.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Osteoporosis bone pain: 1 month. Other diagnoses: 3 months
Other Criteria	For renewal, patient has experienced an objective improvement (i.e. weight gain, increase in lean body mass, or reduction in muscle pain/weakness)

PCSK9 INHIBITOR

Products Affected

- Praluent Subcutaneous Solution Pen-Injector
- Repatha
- Repatha Pushtronex System
- Repatha SureClick

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	<p>For PRALUENT: MUST MEET CRITERIA #1 OR #3. For REPATHA: MUST MEET CRITERIA #1, #2, OR #3. 1. Diagnosis of heterozygous familial hypercholesterolemia (HeFH) confirmed by genotyping OR Simon Broome criteria: Total cholesterol greater than 290mg/dL or LDL cholesterol greater than 190mg/dL, PLUS ONE OF THE FOLLOWING: Tendon xanthomas in patient or 1st degree relative (parent, sibling, child) or 2nd degree relative (grandparent, uncle, aunt) OR DNA-based evidence of LDL receptor mutation, familial defective apo B-100, or PCSK9 mutation. 2a. Myocardial infarction prophylaxis, stroke prophylaxis, and to reduce risk of coronary revascularization in patients with established CVD OR 2b. Diagnosis of homozygous familial hypercholesterolemia (HoFH) confirmed by genotyping OR diagnosis based on the following: a. History of untreated LDL-C greater than 500 mg/dL AND xanthoma before 10 years of age OR b. Documentation of HeFH in both parents. 3. Diagnosis of clinical atherosclerotic cardiovascular disease as defined as one of the following: a. acute coronary syndrome, b. history of myocardial infarction, c. stable/unstable angina, d. coronary or other arterial revascularization, e. stroke, f. transient ischemic stroke, g. peripheral arterial disease presumed to be atherosclerotic region AND MEETS CRITERIA #4, #5, and #6. 4. Provide baseline and current LDL-C. 5. LDL-C greater than or equal to 70mg/dL. 6. Used in combination with maximally tolerated high-intensity statin OR MEETS CRITERIA #7 AND #8. 7. Statin intolerant 8. LDL-C greater than or equal to 70mg/dL. CONTINUING THERAPY: 1. Documented response to Praluent or Repatha, defined as ONE of the following: a. The patient is tolerating medication b. Will continue to be used in combination with maximally tolerated statin (unless statin intolerant).</p>

PA Criteria	Criteria Details
Age Restrictions	Repatha: 13 years of age or older for diagnosis HoFM, Diagnosis CVD or HeFH AND Praluent or Repatha : 18 years of age or older
Prescriber Restrictions	Must be prescribed by, or in consultation with, a cardiologist, endocrinologist, or lipid specialist
Coverage Duration	Initial approval: 8 weeks, Renewal approval: 12 months
Other Criteria	None

PEGYLATED INTERFERON

Products Affected

- Pegasys ProClick

- Pegasys Subcutaneous Solution

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Uncontrolled depression. Autoimmune hepatitis or other autoimmune condition known to be exacerbated by interferon.
Required Medical Information	Chronic Hepatitis C: Criteria will be applied consistent with current AASLD-IDSA guidance OR diagnosis of HBeAg-positive and chronic hepatitis B infection.
Age Restrictions	5 years of age and older. Hepatitis B: 3 years of age and older.
Prescriber Restrictions	Prescribed by or in consultation with a hepatologist, gastroenterologist, oncologist or infectious disease specialist
Coverage Duration	HepC: Initial: 28 wks. Reauth: 20 wks. HepB: 48 weeks
Other Criteria	None

PERJETA

Products Affected

- Perjeta

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of HER2-positive locally advanced, inflammatory, or early stage breast cancer (either greater than 2 cm in diameter or node positive) or metastatic breast cancer AND medication is being used in combination with trastuzumab and docetaxel AND Patient has not received prior anti-HER2 therapy (e.g., trastuzumab) or chemotherapy for metastatic disease AND pregnancy status will be verified prior to initiation of therapy AND females of reproductive potential will be advised of the risks of embryo-fetal death and birth defects, and the need for effective contraception during and after pertuzumab treatment OR Diagnosis of HER2-positive early breast cancer which is at high risk of recurrence AND medication is being used in combination with trastuzumab and chemotherapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

POMALYST

Products Affected

- Pomalyst

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Documentation of ALL of the following: 1. Disease has progressed within 60 days of completion of the last therapy 2. If female of reproductive potential ALL of the below: Two negative pregnancy tests obtained prior to initiating therapy with Pomalyst, monthly negative pregnancy tests during therapy 3. Patient has been counseled about the use of reliable contraception before, during, and 1 month after initiation of therapy with Pomalyst 4. Patient assessment to determine if prophylactic aspirin or antithrombic treatment (warfarin, clopidogrel) will need to be taken to reduce the risk of VTE (embolism, stroke) 5. Registered and certified to be compliant with Pomalyst REMS (Risk Evaluation and Mitigation Strategy) program
Age Restrictions	None
Prescriber Restrictions	Oncologist or hematologist
Coverage Duration	12 months
Other Criteria	A documented diagnosis of multiple myeloma and received at least two prior therapies including lenalidomide (Revlimid) and bortezomib (Velcade)

PROLEUKIN

Products Affected

- Proleukin

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Metastatic Renal Cell Carcinoma or Metastatic Melanoma: Good neurologic or ambulatory performance status (ie, 0 or 1 by Eastern Cooperative Oncology Group, 70-100% by Karnofsky scoring system). Adequate organ function (ie, heart, lungs, kidneys) as determined by all of the following: normal cardiac stress test results, Forced expiratory volume in 1 second (FEV1) greater than 2 L on pulmonary function tests, creatinine concentration 1.5 mg/dL or less.
Age Restrictions	Metastatic Renal Cell Carcinoma or Metastatic Melanoma: 18 years and older
Prescriber Restrictions	Oncologist
Coverage Duration	Metastatic Renal Cell Carcinoma or Metastatic Melanoma: 3 months
Other Criteria	None

PROMACTA

Products Affected

- Promacta

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) Relapsed/refractory chronic immune (idiopathic) thrombocytopenic purpura (ITP) for greater than 6 months AND Baseline platelet count is less than 50,000/mcL AND Degree of thrombocytopenia and clinical condition increase the risk of bleeding AND Patient had an insufficient response, intolerance, contraindication to corticosteroids or immune globulin or inadequate response or contraindication to splenectomy, B)Chronic hepatitis C and patient has thrombocytopenia defined as platelets less than 90,000/mcL for initiation (pre-treatment) of interferon therapy, C) Severe aplastic anemia in patients who have had an insufficient response to immunosuppressive therapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal of ITP, after at least 4 weeks of therapy at the maximum weekly dose (10 mcg/kg) the platelet count increased to a sufficient level to avoid clinically important bleeding. For renewal of Hepatitis C, platelets less than 75,000/mcL for maintenance of optimal interferon-based therapy.

PULMONARY FIBROSIS

Products Affected

- Ofev

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of idiopathic pulmonary fibrosis
Age Restrictions	None
Prescriber Restrictions	Prescriber must be a pulmonologist
Coverage Duration	12 months
Other Criteria	None

PULMOZYME

Products Affected

- Pulmozyme

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations). Part D if patient in long term care (defined by customer location code on claim) otherwise Part B.

QUININE SULFATE

Products Affected

- QuiNINE Sulfate Oral

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Prolongation of QT interval. Glucose-6-phosphate dehydrogenase deficiency. Myasthenia gravis. Known hypersensitivity to mefloquine or quinidine. Optic neuritis. Diagnosis of Blackwater fever
Required Medical Information	Patient has a diagnosis of one of the following: A) uncomplicated Plasmodium falciparum malaria B) uncomplicated Plasmodium vivax malaria C) babesiosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	One month
Other Criteria	None

RADICAVA

Products Affected

- Radicava

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Sulfite hypersensitivity
Required Medical Information	Diagnosis of amyotrophic lateral sclerosis and must meet all of the following: Functionality retained most activities of daily living (defined as scores of 2 points or better on each individual item of the ALS Functional Rating Scale, normal respiratory function defined as percent-predicted forced vital capacity values of percent FVC greater or equal to 80 percent, disease duration of 2 years or less.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in collaboration with a neurologist
Coverage Duration	12 months
Other Criteria	Initial: 6 months. Renewal: 12 months.

RANEXA

Products Affected

- Ranexa

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Hepatic cirrhosis. Pre-existing QT prolongation. Concurrent therapy with a strong CYP3A4 inhibitor. Concurrent therapy with a CYP3A4 inducer.
Required Medical Information	Diagnosis of chronic angina AND patient has tried at least 2 combined anti-anginal therapies such as nitrates, beta-blockers, and calcium channel blockers OR unable to take full doses of conventional angina drugs due to low blood pressure and heart rate.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient had an objective response to therapy

RASUVO

Products Affected

- Rasuvo Subcutaneous Solution Auto-Injector 10 MG/0.2ML, 12.5 MG/0.25ML, 15 MG/0.3ML, 17.5 MG/0.35ML, 20 MG/0.4ML, 22.5 MG/0.45ML, 25 MG/0.5ML, 30 MG/0.6ML, 7.5 MG/0.15ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy or treatment of neoplastic diseases
Required Medical Information	Diagnosis of severe, active rheumatoid arthritis, polyarticular juvenile idiopathic arthritis or severe, recalcitrant, disabling psoriasis AND Failure or clinically significant adverse effects to generic methotrexate injection
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

REGRANEX

Products Affected

- Regranex

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diabetic Neuropathic Ulcers: Diabetic patient with ulcer wound. Treatment will be given in combination with ulcer wound care (eg, debridement, infection control, and/or pressure relief).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Diabetic Neuropathic Ulcers: Maximum 5 months.
Other Criteria	None

REVLIMID

Products Affected

- Revlimid

- Thalomid

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of multiple myeloma and patient has received at least one prior therapy and medication will be used in combination with dexamethasone OR diagnosis of multiple myeloma (maintenance therapy) following autologous hematopoietic stem cell transplantation OR diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes associated with a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities OR diagnosis of mantle cell lymphoma whose disease has relapsed or progressed after two prior therapies, one of which included bortezomib AND patient is enrolled in the Revlimid REMS Program
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

RILUTEK

Products Affected

- Riluzole

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of amyotrophic lateral sclerosis (ALS)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

RUBRACA

Products Affected

- Rubraca

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of deleterious BRCA mutation (germline and/or somatic)-associated epithelial ovarian, fallopian tube, or primary peritoneal cancer and all of the following criteria: 1. BRCA mutation detected by an approved FDA laboratory test, 2. Previous trial/failure with two or more chemotherapy regimens, 3. Used as monotherapy, 4. Agreement of provider to perform a complete blood count (CBC) at baseline and monthly thereafter, 5. Women of reproductive potential must use an effective method of contraception during therapy and for 6 months after the last dose OR Diagnosis of recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer and all of the following: 1. Complete or partial response to platinum-based chemotherapy 2. Used as monotherapy 3. Agreement of provider to perform a complete blood count (CBC) at baseline and monthly thereafter, 4. Women of reproductive potential must use an effective method of contraception during therapy and for 6 months after the last dose.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Hematologist or Oncologist
Coverage Duration	12 months
Other Criteria	None

RYDAPT

Products Affected

- Rydapt

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Angioedema
Required Medical Information	Diagnosis of treatment naive FLT3 mutation-positive acute myelogenous leukemia (AML) and must be used in combination with standard cytarabine and daunorubicin induction and consolidation therapy or diagnosis of systemic mastocytosis.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a oncologist
Coverage Duration	12 months
Other Criteria	None

SABRIL

Products Affected

- Sabril Oral Tablet

- Vigabatrin

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) infantile spasms B) complex partial seizures and patient had an inadequate response to at least one generic first-line agents (carbamazepine, lamotrigine, levetiracetam, oxcarbazepine, valproic acid, divalproex sodium) and at least one adjunctive agent (carbamazepine, clobazam, gabapentin, lamotrigine, levetiracetam, oxcarbazepine, valproic acid, divalproex sodium, topiramate) AND patient and prescriber are enrolled in the SHARE restricted distribution program.
Age Restrictions	Seizures - 10 years of age or older. Infantile spasms - at least one month to 2 years of age
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

SAMSCA

Products Affected

- Samsca

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

SANDOSTATIN

Products Affected

- Octreotide Acetate Injection Solution
100 MCG/ML, 1000 MCG/ML, 200 MCG/ML, 50
MCG/ML, 500 MCG/ML

- SandoSTATIN LAR Depot

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of acromegaly and patient had an inadequate response or cannot be treated with surgical resection, pituitary irradiation, and/or bromocriptine mesylate at maximally tolerated doses OR Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea and flushing episodes OR Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal of acromegaly, IGF-1 level has normalized or improved. For renewal of metastatic carcinoid tumor, patient has improvement in diarrhea and flushing episodes. For renewal of vasoactive intestinal peptide tumor, improvement in diarrhea episodes.

SIGNIFOR

Products Affected

- Signifor

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Acromegaly (initial): Diagnosis of acromegaly AND History of failure to surgery or patient is not a candidate for surgery. Cushing's disease (initial): Diagnosis of endogenous Cushing's disease (i.e., hypercortisolism is not a result of chronic administration of high dose glucocorticoids). Either pituitary surgery has not been curative for the patient OR patient is not a candidate for pituitary surgery. (Reauthorization): Documentation of positive clinical response to Signifor therapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months trial. Reauthorization: 12 months if demonstrated benefit
Other Criteria	Cushing's disease (reauth): a clinically meaningful reduction in 24-hour urinary free cortisol levels or improvement in signs or symptoms of the disease. Acromegaly (reauth): patient's growth hormone (GH) level or insulin-like growth factor 1 (IGF-1) level for age and gender has normalized/improved

SILDENAFIL

Products Affected

- Sildenafil Citrate Intravenous

- Sildenafil Citrate Oral Tablet 20 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

SIMPONI

Products Affected

- Simponi Aria
- Simponi Subcutaneous Solution Auto-Injector
- Simponi Subcutaneous Solution Prefilled Syringe

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	<p>Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: Receiving concurrent therapy with methotrexate (Rheumatrex/Trexall) OR failure, contraindication, or intolerance to methotrexate (Rheumatrex/Trexall). One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), OR for continuation of prior Simponi therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), OR for continuation of prior Simponi therapy. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), OR for continuation of prior Simponi therapy. Ulcerative Colitis (UC) (Initial): Diagnosis of moderately to severely active UC. Patient is corticosteroid dependent (ie, an inability to successfully taper corticosteroids without a return of the symptoms of UC), OR history of failure, contraindication, or intolerance to one of the following: oral aminosalicylate, oral corticosteroid, azathioprine, 6-mercaptopurine. One of the following: Failure, contraindication, or intolerance to Humira (adalimumab), OR for continuation of prior Simponi therapy. All indications (Initial, reauth): Patient is not receiving Simponi in combination with a biologic DMARD [eg, Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Orencia (abatacept)]. Patient is not receiving Simponi in combination with a Janus kinase inhibitor [eg, Xeljanz (tofacitinib)]. For a diagnosis of PsA, Patient is not receiving Simponi in combination with a phosphodiesterase 4 (PDE4) inhibitor [e.g., Otezla (apremilast)].</p>

PA Criteria	Criteria Details
Age Restrictions	None
Prescriber Restrictions	RA, AS (Initial): Prescribed by or in consultation with a rheumatologist. PsA (Initial): Prescribed by or in consultation with a rheumatologist or dermatologist. UC (Initial): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	UC (Initial): 12 weeks. UC (Reauth): 12 months. RA, AS, PsA (Initial, reauth): 12 months
Other Criteria	All indications (Reauth): Documentation of positive clinical response to Simponi therapy.

SOLTAMOX

Products Affected

- Soltamox

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis for use. Documentation of inability to swallow tablet formulation.
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

SOMATULINE DEPOT

Products Affected

- Somatuline Depot

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of acromegaly AND Inadequate response to surgery and/or radiation therapy or patient cannot be treated with surgery and/or radiotherapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient's IGF-1 levels has normalized or improved.

SOMAVERT

Products Affected

- Somavert

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	IV administration of Somavert, concomitant use of Sandostatin or Somatuline.
Required Medical Information	Diagnosis of acromegaly was confirmed by an elevated IGF-1 level or elevated GH level with a glucose tolerance test. Patient has tried and failed at least a 3 month trial of Sandostatin or Somatuline. For renewal, reduction in IGF-1 level from baseline.
Age Restrictions	None
Prescriber Restrictions	Endocrinologist
Coverage Duration	12 months
Other Criteria	None

SPORANOX

Products Affected

- Itraconazole Oral Capsule

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Ventricular dysfunction. Congestive heart failure (CHF). History of CHF. Concurrent therapy with a CYP3A4 inhibitor (e.g., cisapride, lovastatin, methadone, etc.)
Required Medical Information	Patient meets one of the following conditions: A) Diagnosis of systemic fungal infection (e.g., aspergillosis, histoplasmosis, blastomycosis) OR B) Diagnosis of onychomycosis confirmed by one of the following: positive potassium hydroxide (KOH) preparation, culture, or histology and the patient has extensive nail involvement causing significant pain and/or debilitation and Patient has tried or had a contraindication or intolerance to oral terbinafine OR C) Diagnosis of one of the following: tinea corporis (ringworm), tinea cruris (jock itch), tinea pedis (athlete's foot), tinea capitis (scalp ringworm), pityriasis versicolor and the patient is resistant to topical treatment.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Systemic infection: 6 months. Onychomycosis 2 months (fingernail), 3 months (toenail)
Other Criteria	None

SPRYCEL

Products Affected

- Sprycel

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML) that is newly diagnosed in the chronic phase OR Ph+ CML with resistance or intolerance to prior therapy, including imatinib OR Diagnosis of Ph+ acute lymphoblastic leukemia with resistance or intolerance to prior therapy OR Gastrointestinal stromal tumors (GIST) after disease progression on Gleevec (imatinib) or Sutent (sunitinib)
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

STELARA

Products Affected

- Stelara Intravenous

- Stelara Subcutaneous Solution 45 MG/0.5ML
- Stelara Subcutaneous Solution Prefilled Syringe

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Plaque psoriasis (Initial - 45mg/0.5mL): Diagnosis of moderate to severe plaque psoriasis. Plaque psoriasis (Initial - 90mg/1mL): Diagnosis of moderate to severe plaque psoriasis. Patient's weight is greater than 100 kg (220 lbs). Psoriatic arthritis (PsA) (Initial - 45mg/0.5mL): Diagnosis of active PsA. PsA (Initial - 90mg/1mL): Diagnosis of active PsA. Patient's weight is greater than 100 kg (220 lbs). Diagnosis of co-existent moderate to severe psoriasis. Plaque psoriasis (Initial): One of the following: a) History of failure, contraindication, or intolerance (F/C/I) to Enbrel (etanercept) and Humira (adalimumab) OR b) for continuation of prior Stelara therapy. PsA (Initial): One of the following: a) History of failure, contraindication, or intolerance (F/C/I) to Enbrel (etanercept) and Humira (adalimumab) OR b) for continuation of prior Stelara therapy. For Crohn's disease, history of failure, contraindication, or intolerance (F/C/I) to Humira (adalimumab).
Age Restrictions	None
Prescriber Restrictions	Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	All uses (Initial, reauth): 12 months
Other Criteria	Reauthorization (all indications): Documentation of positive clinical response to Stelara therapy. All indications (initial, reauth): Patient is not receiving Stelara in combination with either of the following: Biologic DMARD [eg, Enbrel (etanercept), Humira (adalimumab),

PA Criteria	Criteria Details
	Cimzia (certolizumab), Simponi (golimumab), Otezla (apremilast)] or a Janus Kinase Inhibitor [eg, Xeljanz (tofacitinib)].Patient is not receiving Stelara in combination with a phosphodiesterase 4 (PDE4) inhibitor [e.g., Otezla (apremilast)].

STIVARGA

Products Affected

- Stivarga

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	A documented diagnosis of metastatic colorectal cancer AND documentation of prior therapy with ALL of the following per the indication: 1. (fluoropyrimidine-, oxaliplatin-, and irinotecan)-based chemotherapy 2. bevacizumab (Avastin) 3. panitumumab (Vectibix) OR cetuximab (Erbix) (for KRAS mutation-negative patients only) OR a documented diagnosis of locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) who have been previously treated with imatinib mesylate (Gleevec) and sunitinib malate (Sutent) OR a documented diagnosis of hepatocellular carcinoma in patients previously treated with sorafenib (Nexavar).
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

SUTENT

Products Affected

- Sutent

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced/metastatic renal cell carcinoma OR Diagnosis of gastrointestinal stromal tumors after disease progression on or intolerance to Gleevec OR Diagnosis of progressive, well-differentiated pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease OR Diagnosis of high risk recurrent renal cell carcinoma following nephrectomy, used as adjuvant therapy.
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

SYLATRON

Products Affected

- Sylatron Subcutaneous Kit 200 MCG, 300 MCG, 600 MCG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Autoimmune hepatitis. Hepatic decompensation (Child-Pugh score greater than 6 [Class B or C])
Required Medical Information	Melanoma: Diagnosis of melanoma with microscopic or gross nodal involvement AND The prescribed medication will be used as adjuvant therapy within 84 days of definitive surgical resection, including complete lymphadenectomy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

SYMDEKO

Products Affected

- Symdeko

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis and patient is homozygous for the F508del mutation OR have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor verified by an FDA-cleared CF mutation test
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial-6 months. renewal-12 months.
Other Criteria	None

SYMLIN

Products Affected

- SymlinPen 120 Subcutaneous Solution Pen-Injector

- SymlinPen 60 Subcutaneous Solution Pen-Injector

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Confirmed diagnosis of gastroparesis. Concurrent use of drugs that stimulate gastrointestinal motility. Recurrent severe hypoglycemia requiring assistance during the past 6 months. Presence of hypoglycemia unawareness. Poor compliance with current insulin regimen. Poor compliance with prescribed self-blood glucose monitoring. Hemoglobin A1c level higher than 9%.
Required Medical Information	Diagnosis of type 1 or type 2 diabetes mellitus AND Patient is taking concurrent mealtime insulin therapy (e.g., Humulin, Humalog, Novolin, Novolog)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, the patient had an objective response to therapy.

SYNAGIS

Products Affected

- Synagis

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Patient will use palivizumab for immunoprophylaxis of respiratory syncytial virus (RSV) during the peak months of infection in the patients geographic region AND Patient meets one of the following criteria: A) Infants born at 28 weeks, six days gestation or earlier and who are younger than 12 months of age at the start of the RSV season OR B) Infants born at 29 to 31 weeks, six days gestation and who are younger than six months of age at the start of the RSV season OR C) Infants born at 32 to 34 weeks, six days gestation and who are younger than three months of age at the start of RSV season with at least one of the following risk factors may be dosed until 90 days of age: Child care attendance or Sibling younger than five years of age living in the same household (who is not a multiple birth younger than one year of age) OR D) Infants and children younger than one year of age at the start of RSV season with either congenital abnormalities of the airway or neuromuscular disease that compromises handling of respiratory secretions OR E) Infants and children younger than two years of age with hemodynamically significant congenital heart disease and who have at least one of the following criteria: Receiving medication to control congestive heart failure, Has moderate to severe pulmonary hypertension, or Has cyanotic heart disease OR F) Infants and children younger than two years of age who have received medical therapy (oxygen, bronchodilator, diuretic, or corticosteroid therapy) for chronic lung disease within six months of the start of the RSV season
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	Approve 5 doses based on patient body weight

SYNAREL

Products Affected

- Synarel

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Synarel should not be administered to patients who are hypersensitive to GnRH, GnRH agonist analogues or any of the excipients of SYNAREL, have undiagnosed vaginal bleeding, are pregnant or may become pregnant as major fetal abnormalities were observed in rats (not applicable when used in invitro fertilization programs), are breast feeding.
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

SYNRIBO

Products Affected

- Synribo

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic myelogenous leukemia AND patient has tried and failed or has a contraindication or intolerance to 2 tyrosine kinase inhibitors
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

SYPRINE

Products Affected

- Trientine HCl

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Wilson's disease and intolerance to penicillamine
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

TAFINLAR

Products Affected

- Tafinlar

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	A documented positive BRAF V600E or V600K mutation as detected by an FDA-approved test
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	Tafinlar should not be used in patients with wild-type BRAF melanoma due to the potential risk of tumor promotion in these patients

TAGRISSO

Products Affected

- Tagrisso

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic, non-small cell lung cancer with one of the following- confirmed presence of T790M EGFR tumor mutation positive disease OR confirmed presence of epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R tumor mutations, as detected by an FDA-approved test.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

TARCEVA

Products Affected

- Tarceva

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced, unresectable, or metastatic pancreatic cancer and Tarceva will be used in combination with gemcitabine OR Diagnosis of locally advanced or metastatic (stage III or IV) non-small cell lung cancer with one of the following: A) failure with at least one prior chemotherapy regimen and Tarceva will be used as monotherapy, or B) no evidence of disease progression after four cycles of first-line platinum-based chemotherapy and Tarceva will be used as maintenance treatment and Tarceva will be used as monotherapy, or C) Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutation as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

TASIGNA

Products Affected

- Tasigna

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Long QT syndrome. Uncorrected hypokalemia. Uncorrected hypomagnesemia. Concomitant use with a drug known to prolong the QT interval or strong cytochrome P450 3A4 inhibitors
Required Medical Information	Diagnosis of newly diagnosed Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML) in the chronic phase OR Diagnosis of Ph+ CML with resistance or intolerance to prior therapy that include imatinib.
Age Restrictions	1 year of age and older
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

TAZORAC

Products Affected

- Tazarotene External

- Tazorac External Cream 0.05 %
- Tazorac External Gel

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of acne vulgaris and patient has tried an adequate trial with at least one other topical acne product (e.g., benzoyl peroxide, salicylic acid, clindamycin, erythromycin, adapalene, azelaic acid, and/or tretinoin) OR Diagnosis of stable moderate to severe plaque psoriasis and 20% or less body surface area involvement and patient has a contraindication or tried adequate trial with at least one other topical psoriasis product (e.g., medium to high potency corticosteroid and/or vitamin D analogs) AND females of child-bearing potential are using adequate birth control measures during therapy.
Age Restrictions	12 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

TECENTRIQ

Products Affected

- Tecentriq

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic urothelial carcinoma. One of the following: A) History of disease progression during or following platinum-containing chemotherapy, OR B) History of disease progression within 12 months of neoadjuvant or adjuvant treatment with a platinum-containing chemotherapy.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

TESTOSTERONE

Products Affected

- Testosterone Cypionate Intramuscular Solution 100 MG/ML, 200 MG/ML
- Testosterone Enanthate Intramuscular Solution
- Testosterone Transdermal Gel 10 MG/ACT (2%), 12.5 MG/ACT (1%), 50 MG/5GM (1%)
- Testosterone Transdermal Solution

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Carcinoma of the breast. Known or suspected carcinoma of the prostate.
Required Medical Information	Diagnosis of hypogonadism (primary or hypogonadotropic) AND patient is male AND patient's serum testosterone (total or free) value and the laboratory reference value range reported by laboratory service AND diagnosis has been confirmed by a low-for-age serum testosterone (total or free) level defined by the normal laboratory reference value
Age Restrictions	12 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, patient experienced an objective response to therapy.

TETRABENAZINE

Products Affected

- Tetrabenazine

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Actively suicidal. Untreated or inadequately treated depression. Impaired hepatic function. Concomitant use of monoamine oxidase inhibitors. Concomitant use of reserpine or within 20 days of discontinuing reserpine.
Required Medical Information	Diagnosis of chorea associated with Huntington's disease AND any medication possibly contributing to the underlying symptoms of chorea has been discontinued (e.g., antipsychotics, metoclopramide, amphetamines, methylphenidate, dopamine agonists, etc.) unless cessation would be detrimental to the underlying condition AND patient has been genotyped to CYP2D6 to determine whether the patient is a poor, intermediate or extensive metabolizer.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Dosing will be approved per the FDA labeling based on CYP2D6 testing. For renewal, patient had an objective response to therapy.

TOPICAL RETINOIDS

Products Affected

- Adapalene External Cream
- Adapalene External Gel

- Tretinoin External Cream
- Tretinoin External Gel 0.01 %, 0.025 %

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate acne vulgaris
Age Restrictions	PA applies to patients older than 26 years of age
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, medication has been effective in treating the patient's condition.

TRACLEER

Products Affected

- Tracleer

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Receiving concomitant cyclosporine A or glyburide therapy. Aminotransferase elevations are accompanied by signs or symptoms of liver dysfunction or injury or increases in bilirubin at least 2 times the upper limit of normal.
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND Patient has WHO Group I PAH AND Patient has New York Heart Association (NYHA) Functional Class II-IV AND pregnancy must be excluded prior to the start of therapy and will be prevented thereafter with two forms of reliable contraception in female patients of reproductive potential.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months - initial. 12 months - renewal
Other Criteria	Liver aminotransferases will be measured prior to initiation of treatment and then monthly.

TRELSTAR

Products Affected

- Trelstar Mixject

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of FDA approved indications not otherwise excluded from Part D AND palliative treatment of advanced prostate cancer, central precocious puberty, endometrial hyperplasia, endometriosis, fibrocystic disease of breast, uterine leiomyoma
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

TYKERB

Products Affected

- Tykerb

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) AND a) the medication will be used in combination with Xeloda in a patient with advanced or metastatic disease and the patient has received prior therapy including an anthracycline, a taxane, and trastuzumab or b) The medication will be used in combination with Femara for the treatment of a postmenopausal woman with hormone receptor-positive metastatic disease for whom hormonal therapy is indicated.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

TYMLOS

Products Affected

- Tymlos

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Patients at increased risk of osteogenic sarcoma.
Required Medical Information	Diagnosis of osteoporosis in post-menopausal women at high risk for fracture. Member must have failed therapy with a bisphosphonate (defined by a fracture while on therapy or worsening bone density) unless such a trial is shown to be inappropriate or contraindicated (i.e., presence of severe osteoporosis [T-scores -3.0 or worse in lumbar spine, femoral neck, or total hip region], history of major osteoporotic fracture, presence of renal insufficiency, etc) AND member has at least one of the following: T-score equal to or worse than -2.5 in the lumbar spine, femoral neck, or total hip region OR a FRAX calculator based 10-year risk of at least 20% for a major osteoporotic fracture (spine, shoulder, hip, or wrist), or a 10-year risk of at least 3% for a hip fracture OR presence or history of osteoporotic fracture.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 12 months. Reauth: Treatment duration has not exceeded 24 months during patient lifetime.
Other Criteria	None

TYSABRI

Products Affected

- Tysabri

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	History of progressive multifocal leukoencephalopathy.
Required Medical Information	MULTIPLE SCLEROSIS: TRIAL OF AN INTERFERON OR COPAXONE. CROHN'S DISEASE: TRIAL OF A TNF-ALPHA INHIBITOR. RENEWAL: CROHN'S: PATIENT IS NOT ON CONCOMITANT CORTICOSTEROID TREATMENT AFTER 6 MONTHS ON NATALIZUMAB, OR HAS NOT RECEIVED MORE THAN 3 MONTHS OF A CORTICOSTEROID WITHIN THE PAST 12 MONTHS.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	MULTIPLE SCLEROSIS:12 MONTHS. CROHN'S DISEASE: 6 MONTHS. RENEWAL: CROHN'S: 12 MONTHS.
Other Criteria	Patient and physician are registered in the TOUCH prescribing program. For renewal, patient had an objective response to therapy (e.g., decreased flare).

UPTRAVI

Products Affected

- Upravi

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group 1) confirmed by right heart catheterization AND patient has tried and had an insufficient response to at least one other PAH agent (e.g., sildenafil) therapy and one other ERA agent (e.g. letairis, opsumit, tracleer).
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

VALCHLOR

Products Affected

- Valchlor

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of mycosis fungoides-type cutaneous T-cell lymphoma AND patient has early stage disease (defined as Stage 1A or 1B) AND patient has received prior skin-directed therapy (e.g., very high potency class I topical corticosteroids for at least 3 months (i.e. clobetasol, diflorasone, halobetasol, augmented betamethasone dipropionate), phototherapy, topical nitrogen mustard, or a topical retinoid (e.g., bexarotene)).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

VARIZIG

Products Affected

- VariZIG Intramuscular Solution

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	History of hypersensitivity (including anaphylaxis or severe systemic reaction) to immune globulin or any component of the preparation. Severe thrombocytopenia or coagulation disorder where IM injections are contraindicated.
Required Medical Information	To be used for post-exposure varicella infection prophylaxis to reduce varicella severity in high-risk patients defined as premature neonates, neonates and infants less than 1 year old, pregnant women, newborns of women with varicella shortly before or after delivery, immunocompromised children and adults without a past history of varicella unless the patient is undergoing a bone marrow transplantation, and adults without evidence of immunity.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None

VECTIBIX

Products Affected

- Vectibix Intravenous Solution 100 MG/5ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of epidermal growth factor receptor (EGFR)-expressing, metastatic colorectal carcinoma with disease progression following fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy regimens OR Diagnosis of wild-type KRAS metastatic colorectal cancer and used in combination with FOLFOX therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Subject to B vs D

VELCADE

Products Affected

- Bortezomib

- Velcade Injection

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Hypersensitivity to bortezomib, boron, or mannitol. Intrathecal administration.
Required Medical Information	Diagnosis of mantle cell lymphoma OR multiple myeloma and at least 1 prior therapy
Age Restrictions	None
Prescriber Restrictions	Oncologist or hematologist
Coverage Duration	12 months
Other Criteria	All uses: for continuation of therapy

VENCLEXTA

Products Affected

- Venclexta

- Venclexta Starting Pack

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic lymphocytic leukemia (CLL) OR small lymphocytic lymphoma, with or without 17p deletion and patient has had at least 1 prior therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

VENTAVIS

Products Affected

- Ventavis

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Congestive heart failure due to severe left ventricular systolic dysfunction.
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND Patient has WHO Group I PAH AND Patient has New York Heart Association (NYHA) Functional Class III or IV.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial - 6 months. Renewal - 12 months
Other Criteria	Part D if patient in long term care (defined by customer location code on claim) otherwise Part B.

VERZENIO

Products Affected

- Verzenio

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	BREAST CANCER (1) Patient must have a diagnosis of advanced or metastatic breast cancer AND (2a) must be used in combination with fulvestrant for the treatment of disease progression following endocrine therapy OR (2b) used as monotherapy for treatment of disease progression following endocrine therapy and patient has already received at least one prior chemotherapy regimen of Ibrance or Kisqali OR (2c) used as initial endocrine-based treatment in combination with an aromatase inhibitor AND (3) disease is hormone receptor positive AND human epidermal growth factor 2 (HER2)-negative
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of therapy

VOTRIENT

Products Affected

- Votrient

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced/metastatic renal cell carcinoma OR Diagnosis of advanced soft tissue sarcoma and patient received at least one prior chemotherapy (e.g., doxorubicin, dacarbazine, ifosfamide, epirubicin, docetaxel, or vinorelbine).
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

VPRIV

Products Affected

- Vpriv

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of type 1 Gaucher disease
Age Restrictions	4 years of age or older
Prescriber Restrictions	None
Coverage Duration	3 months - initial. 12 months - renewal
Other Criteria	None

VYXEOS

Products Affected

- Vyxeos Intravenous Suspension Reconstituted 44-100 MG

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of therapy related acute myeloid leukemia or acute myeloid leukemia with myelodysplasia related changes. If the patient has the diagnosis of therapy related acute myeloid leukemia, it must be newly diagnosed.
Age Restrictions	age 18 years and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	BvD determination per CMS guidelines

XALKORI

Products Affected

- Xalkori

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic (stage III or IV) non-small cell lung cancer AND patient has non-squamous cell histology AND Patient has anaplastic lymphoma kinase (ALK)-positive disease as detected with an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility or are ROS1-positive
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

XELJANZ

Products Affected

- Xeljanz

- Xeljanz XR

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Psoriatic arthritis (PsA) or Rheumatoid arthritis (RA) or Ulcerative Colitis (Initial): Diagnosis of psoriatic arthritis or moderately to severely active RA and an inadequate response or intolerance to methotrexate. One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab) (for Ulcerative Colitis only Humira will be required), OR patient has a documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-IV-TR 300.29 for specific phobia diagnostic criteria), OR for continuation of prior tofacitinib therapy. Patient is not receiving tofacitinib in combination with a biologic DMARD [eg, Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab)]. Patient is not receiving tofacitinib in combination with a potent immunosuppressant (eg, azathioprine, cyclosporine).
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist or gastroenterologist
Coverage Duration	12 months
Other Criteria	Documentation of positive clinical response to tofacitinib therapy. Patient is not receiving tofacitinib in combination with a biologic DMARD [eg, Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab)]. Patient is not receiving tofacitinib in combination with a potent immunosuppressant (eg, azathioprine, cyclosporine).

XGEVA

Products Affected

- Xgeva

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

XOLAIR

Products Affected

- Xolair

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Documentation of the following: A) moderate to severe chronic idiopathic urticaria and has remained symptomatic despite at least 2 weeks of H1 antihistamine therapy OR intolerance or contraindication of H1 antihistamine therapy OR B) Mod-severe persistent asthma (NHLBI definition) meeting all the following criteria: Evidence of reversible disease (12% or greater improvement in FEV1 with at least a 200ml increase or 20% or greater improvement in PEF as a result of a short-acting bronchodilator challenge). Evidence of specific allergic sensitivity to a perennial aeroallergen (+ skin test or in vitro test). Failure of an adequate trial of standard therapy as defined by a trial of at least a 3 month course of high-dose inhaled corticosteroids and long-acting beta2-agonists OR maximally tolerated doses of standard therapy OR intolerance or contraindication to standard therapy. Extended approval for 6 months if demonstrated benefit, meeting at least 2 of the following criteria: PEF improvement (12% or greater from baseline (prior to start of Xolair)), OR FEV1 improvement (12% or greater from baseline (prior to start of Xolair)), OR reduction in symptoms (wheezing, sob, cough, chest tightness), OR reduction in systemic corticosteroids and rescue drug use, OR reduction of asthma-related hospitalizations and other medical contacts.
Age Restrictions	6 years of age or older
Prescriber Restrictions	Allergist, immunologist, pulmonologist or dermatologist
Coverage Duration	Initial: 6 months trial. Extended approval: 6 months if demonstrated benefit
Other Criteria	If this medication is administered by a physician incident to a physicians visit this would be covered by Medicare Part B

XTANDI

Products Affected

- Xtandi

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Metastatic castration-resistant prostate cancer (mCRPC): Diagnosis of mCRPC. History of failure, contraindication or intolerance to Zytiga.
Age Restrictions	None
Prescriber Restrictions	Prescribed or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

XURIDEN

Products Affected

- Xuriden

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of Hereditary orotic aciduria
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a specialist that treats metabolic defects
Coverage Duration	12 months
Other Criteria	None

XYREM

Products Affected

- Xyrem

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Concomitant treatment with sedative hypnotic agents. Succinic semialdehyde dehydrogenase deficiency.
Required Medical Information	Diagnosis of narcolepsy with excessive daytime sleepiness, cataplexy or both and for patients with excessive daytime sleepiness, patient has had a previous trial with or has a contraindication, intolerance, or allergy to Provigil or Nuvigil.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	For renewal, the patient had a positive response to the medication (increased sleep quality for patients with narcolepsy)

YERVOY

Products Affected

- Yervoy Intravenous Solution 50 MG/10ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic melanoma AND If the request is for re-induction, the patient had no significant toxicity with the prior course of Yervoy AND the patient experienced progression after having stable disease for longer than three months or relapse after having a clinical response to therapy AND the prescriber is aware of the Yervoy REMS program OR used as adjuvant therapy for the diagnosis of cutaneous melanoma in patients with pathologic involvement of regional lymph nodes of more than 1mm who have undergone complete resection, including total lymphadenectomy OR diagnosis of advanced renal cell carcinoma and used in combination with Opdivo
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	16 weeks, 12 months when used as adjuvant therapy
Other Criteria	Authorization will be for 4 doses for unresectable or metastatic melanoma

YONDELIS

Products Affected

- Yondelis

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis and lab values: ANC, platelet count, serum creatine phosphokinase, serum creatinine, liver function tests, and left ventricular ejection fraction.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Must be prescribed by an oncologist
Coverage Duration	12 months
Other Criteria	None

YONSA

Products Affected

- Yonsa

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic castration-resistant prostate cancer, and used in combination with methylprednisolone AND Documented history of trial with, inadequate treatment response, adverse event, or contraindication to Zytiga
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None

ZALTRAP

Products Affected

- Zaltrap Intravenous Solution 100 MG/4ML

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Severe hemorrhage, development of gastrointestinal perforation, compromised wound healing
Required Medical Information	Diagnosis of metastatic colorectal cancer AND will be used in combination with 5-fluorouracil, leucovorin, and irinotecan (FOLFIRI) AND disease is resistant to or has progressed following an oxaliplatin-containing regimen
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Patient will be monitored for signs and symptoms of gastrointestinal bleeding and other severe bleeding. Therapy will be suspended for at least 4 weeks prior to elective surgery and not resumed for at least 4 weeks following major surgery and until the wound is fully healed.

ZAVESCA

Products Affected

- Miglustat

- Zavesca

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate type 1 Gaucher disease and patient is not a candidate for enzyme replacement therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ZEJULA

Products Affected

- Zejula

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of recurrent epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer and patient had a complete or partial response to platinum-based chemotherapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a oncologist or gynecologist
Coverage Duration	12 months
Other Criteria	None

ZELBORAF

Products Affected

- Zelboraf

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic melanoma and patient has positive BRAF-V600E mutation documented by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility.
Age Restrictions	None
Prescriber Restrictions	Oncologist
Coverage Duration	12 months
Other Criteria	None

ZOLEDRONIC ACID

- Products Affected**
- Zoledronic Acid Intravenous Concentrate MG/100ML
 - Zoledronic Acid Intravenous Solution 5

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	Current treatment with Reclast.
Required Medical Information	Diagnosis of hypercalcemia of malignancy and has a corrected calcium greater than or equal to 12 mg/dL OR Diagnosis of multiple myeloma and associated bone disease (e.g., osteolytic bone lesions, bone metastases, osteopenia, etc.) OR Diagnosis of a solid tumor (e.g., breast cancer, prostate cancer that has progressed after at least one hormonal therapy (i.e. antiandrogen [bicalutamide, flutamide, nilutamide], LHRH agonist [leuprolide, goserelin], LHRH antagonists [degarelix]), kidney cancer, non-small cell lung cancer, or thyroid cancer) and patient has bone metastases and medication will be used in conjunction with standard antineoplastic therapy and medication is used for the prevention of skeletal-related events (e.g. spinal cord compression, hypercalcemia, bone pain or lesions requiring radiation or surgery) AND Patient has tried and had an inadequate response or has a contraindication/intolerance to pamidronate
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ZOLINZA

Products Affected

- Zolinza

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma AND progressive, persistent or recurrent disease or patient is not a candidates for or following 2 systemic therapies (e.g., bexarotene, romidepsin, etc.)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ZORTRESS

Products Affected

- Zortress

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Medication is being used for: A) Prevention of kidney transplant organ rejection AND patient is at low-to-moderate immunologic risk AND member is prescribed concurrent therapy with reduced doses of cyclosporine and corticosteroids, or B) Prevention of liver transplant organ rejection AND 30 or more days have passed since the transplant procedure AND the member is prescribed concurrent therapy with reduced doses of tacrolimus and corticosteroids
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescriber is experienced in immunosuppressive therapy and management of transplant patients.
Coverage Duration	12 months
Other Criteria	Part B if transplant covered by Medicare. otherwise Part D

ZYDELIG

Products Affected

- Zydelig

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	The patient has one of the following diagnoses: A) chronic lymphocytic leukemia AND The medication will be used in combination with rituximab AND The patient has relapsed on at least one prior therapy (e.g., purine analogues [fludarabine, pentostatin, cladribine], alkylating agents [chlorambucil, cyclophosphamide], or monoclonal antibodies [rituximab]) AND the patient does not have any co-morbidities that prevents the use of cytotoxic chemotherapy (i.e. severe neutropenia or thrombocytopenia, creatinine clearance less than 60 mL/minute), B) follicular lymphoma AND the patient has relapsed on at least two prior systemic therapies (e.g., rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine]), or C) small lymphocytic lymphoma AND The patient has relapsed on at least two prior systemic therapies(e.g., rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine]).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ZYKADIA

Products Affected

- Zykadia

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

ZYTIGA

Products Affected

- Zytiga

PA Criteria	Criteria Details
Covered Uses	All medically accepted indications not otherwise excluded from Part D
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic prostate cancer AND Patient has castration-resistant disease (defined by tumor growth/disease progression, risk in PSA levels, new metastases) OR high-risk castration-sensitive prostate cancer AND Zytiga will be used in combination with prednisone.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None

PART B VERSUS PART D

Products Affected

- Abelcet Intravenous Suspension 5 MG/ML
- Abraxane Intravenous Suspension Reconstituted 100 MG
- Acetylcysteine Inhalation Solution 10 %, 20 %
- Acyclovir Sodium Intravenous Solution 50 MG/ML
- Adrucil Intravenous Solution 500 MG/10ML
- Albuterol Sulfate Inhalation Nebulization Solution (2.5 MG/3ML) 0.083%, (5 MG/ML) 0.5%, 0.63 MG/3ML, 1.25 MG/3ML
- Alimta Intravenous Solution Reconstituted 100 MG, 500 MG
- AmBisome Intravenous Suspension Reconstituted 50 MG
- Amikacin Sulfate Injection Solution 500 MG/2ML
- Aminosyn II Intravenous Solution 10 %, 8.5 %
- Aminosyn II/Electrolytes Intravenous Solution 8.5 %
- Aminosyn/Electrolytes Intravenous Solution 7 %, 8.5 %
- Aminosyn-HBC Intravenous Solution 7 %
- Aminosyn-PF Intravenous Solution 10 %, 7 %
- Aminosyn-RF Intravenous Solution 5.2 %
- Amiodarone HCl Intravenous Solution 150 MG/3ML
- Amphotericin B Injection Solution Reconstituted 50 MG
- Ampicillin-Sulbactam Sodium Injection Solution Reconstituted 15 (10-5) GM
- Aprepitant Oral Capsule 125 MG, 40 MG, 80 & 125 MG, 80 MG
- Argatroban Intravenous Solution 125 MG/125ML, 250 MG/2.5ML
- Arranon Intravenous Solution 5 MG/ML
- Astagraf XL Oral Capsule Extended Release 24 Hour 0.5 MG, 1 MG, 5 MG
- Atgam Intravenous Injectable 50 MG/ML
- AzaCITIDine Injection Suspension Reconstituted 100 MG
- Azactam Injection Solution Reconstituted 1 GM, 2 GM
- Azasan Oral Tablet 100 MG, 75 MG
- AzaTHIOprine Oral Tablet 50 MG
- AzaTHIOprine Sodium Injection Solution Reconstituted 100 MG
- Azithromycin Intravenous Solution Reconstituted 500 MG
- BiCNU Intravenous Solution Reconstituted 100 MG
- Bleomycin Sulfate Injection Solution Reconstituted 30 UNIT
- Budesonide Inhalation Suspension 0.25 MG/2ML, 0.5 MG/2ML, 1 MG/2ML
- Busulfan Intravenous Solution 6 MG/ML
- Calcitonin (Salmon) Nasal Solution 200 UNIT/ACT
- Calcitriol Intravenous Solution 1 MCG/ML
- Calcitriol Oral Capsule 0.25 MCG, 0.5 MCG
- Calcitriol Oral Solution 1 MCG/ML
- CARBOplatin Intravenous Solution 150 MG/15ML
- Caspofungin Acetate Intravenous Solution Reconstituted 50 MG, 70 MG
- CefOXitin Sodium Intravenous Solution Reconstituted 1 GM, 2 GM
- CefTRIAXone Sodium Injection Solution Reconstituted 1 GM, 2 GM, 250 MG, 500 MG
- CefTRIAXone Sodium Intravenous Solution Reconstituted 10 GM
- Cerezyme Intravenous Solution Reconstituted 400 UNIT
- Chloramphenicol Sod Succinate Intravenous Solution Reconstituted 1 GM
- Cidofovir Intravenous Solution 75 MG/ML

- Ciprofloxacin in D5W Intravenous Solution 200 MG/100ML
- CISplatin Intravenous Solution 50 MG/50ML
- Cladribine Intravenous Solution 10 MG/10ML
- Clindamycin Phosphate Injection Solution 300 MG/2ML, 900 MG/6ML
- Clinimix E/Dextrose (2.75/10) Intravenous Solution 2.75 %
- Clinimix E/Dextrose (2.75/5) Intravenous Solution 2.75 %
- Clinimix E/Dextrose (4.25/10) Intravenous Solution 4.25 %
- Clinimix E/Dextrose (4.25/25) Intravenous Solution 4.25 %
- Clinimix E/Dextrose (4.25/5) Intravenous Solution 4.25 %
- Clinimix E/Dextrose (5/15) Intravenous Solution 5 %
- Clinimix E/Dextrose (5/20) Intravenous Solution 5 %
- Clinimix E/Dextrose (5/25) Intravenous Solution 5 %
- Clinimix/Dextrose (2.75/5) Intravenous Solution 2.75 %
- Clinimix/Dextrose (4.25/10) Intravenous Solution 4.25 %
- Clinimix/Dextrose (4.25/20) Intravenous Solution 4.25 %
- Clinimix/Dextrose (4.25/25) Intravenous Solution 4.25 %
- Clinimix/Dextrose (4.25/5) Intravenous Solution 4.25 %
- Clinimix/Dextrose (5/15) Intravenous Solution 5 %
- Clinimix/Dextrose (5/20) Intravenous Solution 5 %
- Clinimix/Dextrose (5/25) Intravenous Solution 5 %
- Clinisol SF Intravenous Solution 15 %
- Clofarabine Intravenous Solution 1 MG/ML
- Cromolyn Sodium Inhalation Nebulization Solution 20 MG/2ML
- Cyclophosphamide Oral Capsule 25 MG, 50 MG
- CycloSPORINE Intravenous Solution 50 MG/ML
- CycloSPORINE Modified Oral Capsule 100 MG, 25 MG, 50 MG
- CycloSPORINE Modified Oral Solution 100 MG/ML
- CycloSPORINE Oral Capsule 100 MG, 25 MG
- Cyramza Intravenous Solution 100 MG/10ML, 500 MG/50ML
- Cytarabine (PF) Injection Solution 100 MG/ML
- Cytarabine Injection Solution 20 MG/ML
- Dacarbazine Intravenous Solution Reconstituted 200 MG
- DACTINomycin Intravenous Solution Reconstituted 0.5 MG
- DAPTOMycin Intravenous Solution Reconstituted 500 MG
- DAUNOrubicin HCl Intravenous Injectable 5 MG/ML
- Decitabine Intravenous Solution Reconstituted 50 MG
- Depo-Provera Intramuscular Suspension 400 MG/ML
- Dexamethasone Sodium Phosphate Injection Solution 10 MG/ML
- Dexrazoxane Intravenous Solution Reconstituted 250 MG
- Dextrose in Lactated Ringers Intravenous Solution 5 %
- Dextrose Intravenous Solution 10 %, 5 %
- Dextrose-NaCl Intravenous Solution 10-0.2 %, 10-0.45 %, 2.5-0.45 %, 5-0.2 %, 5-0.225 %, 5-0.33 %, 5-0.45 %, 5-0.9 %
- Diltiazem HCl Intravenous Solution 50 MG/10ML
- Diltiazem HCl Intravenous Solution Reconstituted 100 MG

- Diphtheria-Tetanus Toxoids DT Intramuscular Suspension 25-5 LFU/0.5ML
- DOCEtaxel Intravenous Concentrate 80 MG/4ML
- DOCEtaxel Intravenous Solution 160 MG/16ML
- DOXOrubicin HCl Intravenous Solution 2 MG/ML
- DOXOrubicin HCl Liposomal Intravenous Injectable 2 MG/ML
- Doxy 100 Intravenous Solution Reconstituted 100 MG
- Dronabinol Oral Capsule 10 MG, 2.5 MG, 5 MG
- Elitek Intravenous Solution Reconstituted 1.5 MG, 7.5 MG
- Emend Intravenous Solution Reconstituted 150 MG
- Emend Oral Suspension Reconstituted 125 MG
- Engerix-B Injection Suspension 10 MCG/0.5ML, 20 MCG/ML
- Envarsus XR Oral Tablet Extended Release 24 Hour 0.75 MG, 1 MG, 4 MG
- Epirubicin HCl Intravenous Solution 200 MG/100ML
- Eraxis Intravenous Solution Reconstituted 100 MG, 50 MG
- Erythrocine Lactobionate Intravenous Solution Reconstituted 500 MG
- Esomeprazole Sodium Intravenous Solution Reconstituted 20 MG, 40 MG
- Etopophos Intravenous Solution Reconstituted 100 MG
- Etoposide Intravenous Solution 100 MG/5ML
- Famotidine Intravenous Solution 20 MG/2ML
- Famotidine Premixed Intravenous Solution 20-0.9 MG/50ML-%
- Faslodex Intramuscular Solution 250 MG/5ML
- Flebogamma DIF Intravenous Solution 5 GM/50ML
- Fluconazole in Sodium Chloride Intravenous Solution 200-0.9 MG/100ML-%, 400-0.9 MG/200ML-%
- Fludarabine Phosphate Intravenous Solution Reconstituted 50 MG
- Fluorouracil Intravenous Solution 5 GM/100ML
- Folutyn Intravenous Solution 40 MG/2ML
- FreAmine HBC Intravenous Solution 6.9 %
- GamaSTAN S/D Intramuscular Injectable (10ML), (2ML)
- Gammagard Injection Solution 2.5 GM/25ML
- Gammagard S/D Less IgA Intravenous Solution Reconstituted 10 GM, 5 GM
- Gammaplex Intravenous Solution 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 5 GM/50ML
- Ganciclovir Sodium Intravenous Solution Reconstituted 500 MG
- Gemcitabine HCl Intravenous Solution Reconstituted 1 GM
- Gengraf Oral Capsule 100 MG, 25 MG
- Gengraf Oral Solution 100 MG/ML
- Granisetron HCl Intravenous Solution 0.1 MG/ML, 1 MG/ML, 4 MG/4ML
- Granisetron HCl Oral Tablet 1 MG
- Heparin (Porcine) in D5W Intravenous Solution 40-5 UNIT/ML-%, 50-5 UNIT/ML-%
- Heparin Sod (Porcine) in D5W Intravenous Solution 100 UNIT/ML
- Hepatamine Intravenous Solution 8 %
- Ibandronate Sodium Intravenous Solution 3 MG/3ML
- IDArubicin HCl Intravenous Solution 10 MG/10ML
- Ifosfamide Intravenous Solution Reconstituted 1 GM
- Imipenem-Cilastatin Intravenous Solution Reconstituted 250 MG, 500 MG
- Imogam Rabies-HT Injection Solution 300 UNIT/2ML

- Intralipid Intravenous Emulsion 20 %, 30 %
- Ipratropium Bromide Inhalation Solution 0.02 %
- Ipratropium-Albuterol Inhalation Solution 0.5-2.5 (3) MG/3ML
- Irinotecan HCl Intravenous Solution 100 MG/5ML
- Isolyte-S Intravenous Solution
- Jevtana Intravenous Solution 60 MG/1.5ML
- KCl in Dextrose-NaCl Intravenous Solution 10-5-0.45 MEQ/L-%-%, 20-5-0.2 MEQ/L-%-%, 20-5-0.33 MEQ/L-%-%, 20-5-0.45 MEQ/L-%-%, 20-5-0.9 MEQ/L-%-%, 30-5-0.45 MEQ/L-%-%, 40-5-0.45 MEQ/L-%-%, 40-5-0.9 MEQ/L-%-%
- KCl-Lactated Ringers-D5W Intravenous Solution 20 MEQ/L
- Kepivance Intravenous Solution Reconstituted 6.25 MG
- Kyprolis Intravenous Solution Reconstituted 30 MG, 60 MG
- Labetalol HCl Intravenous Solution 5 MG/ML
- Lactated Ringers Intravenous Solution
- Leucovorin Calcium Injection Solution Reconstituted 100 MG, 350 MG
- LevETIRAcetam in NaCl Intravenous Solution 1000 MG/100ML, 1500 MG/100ML, 500 MG/100ML
- LevETIRAcetam Intravenous Solution 500 MG/5ML
- Levofloxacin in D5W Intravenous Solution 500 MG/100ML, 750 MG/150ML
- Levofloxacin Intravenous Solution 25 MG/ML
- Levoleucovorin Calcium Intravenous Solution 175 MG/17.5ML
- LEVOleucovorin Calcium Intravenous Solution Reconstituted 50 MG
- Lumizyme Intravenous Solution Reconstituted 50 MG
- Melphalan HCl Intravenous Solution Reconstituted 50 MG
- Meropenem Intravenous Solution Reconstituted 1 GM, 500 MG
- Mesna Intravenous Solution 100 MG/ML
- Methotrexate Oral Tablet 2.5 MG
- Methotrexate Sodium (PF) Injection Solution 250 MG/10ML, 50 MG/2ML
- Methotrexate Sodium Injection Solution 250 MG/10ML
- Methotrexate Sodium Injection Solution Reconstituted 1 GM
- MethylPREDNISolone Sodium Succ Injection Solution Reconstituted 1000 MG, 125 MG, 40 MG
- Metoprolol Tartrate Intravenous Solution 5 MG/5ML
- Metoprolol Tartrate Intravenous Solution Cartridge 5 MG/5ML
- MetroNIDAZOLE in NaCl Intravenous Solution 500-0.79 MG/100ML-%
- Miacalcin Injection Solution 200 UNIT/ML
- MitoMYcin Intravenous Solution Reconstituted 20 MG, 40 MG, 5 MG
- Mitoxantrone HCl Intravenous Concentrate 25 MG/12.5ML
- Moxifloxacin HCl in NaCl Intravenous Solution 400 MG/250ML
- Mustargen Injection Solution Reconstituted 10 MG
- Mycamine Intravenous Solution Reconstituted 100 MG, 50 MG
- Mycophenolate Mofetil HCl Intravenous Solution Reconstituted 500 MG
- Mycophenolate Mofetil Oral Capsule 250 MG
- Mycophenolate Mofetil Oral Suspension Reconstituted 200 MG/ML
- Mycophenolate Mofetil Oral Tablet 500 MG
- Mycophenolate Sodium Oral Tablet Delayed Release 180 MG, 360 MG

- Nebupent Inhalation Solution Reconstituted 300 MG
- Nephroamine Intravenous Solution 5.4 %
- Nitroglycerin Intravenous Solution 5 MG/ML
- Normosol-M in D5W Intravenous Solution
- Normosol-R in D5W Intravenous Solution
- Normosol-R pH 7.4 Intravenous Solution
- Nutrilipid Intravenous Emulsion 20 %
- Ondansetron HCl Injection Solution 4 MG/2ML, 4 MG/2ML (2ML SYRINGE)
- Ondansetron HCl Oral Solution 4 MG/5ML
- Ondansetron HCl Oral Tablet 24 MG, 4 MG, 8 MG
- Ondansetron Oral Tablet Dispersible 4 MG, 8 MG
- Oxaliplatin Intravenous Solution 100 MG/20ML
- Oxaliplatin Intravenous Solution Reconstituted 100 MG
- PACLitaxel Intravenous Concentrate 100 MG/16.7ML
- Pamidronate Disodium Intravenous Solution 30 MG/10ML, 6 MG/ML, 90 MG/10ML
- Pantoprazole Sodium Intravenous Solution Reconstituted 40 MG
- Paricalcitol Intravenous Solution 2 MCG/ML, 5 MCG/ML
- Paricalcitol Oral Capsule 1 MCG, 2 MCG, 4 MCG
- Piperacillin Sod-Tazobactam So Intravenous Solution Reconstituted 2.25 (2-0.25) GM, 3.375 (3-0.375) GM, 4.5 (4-0.5) GM, 40.5 (36-4.5) GM
- Plasma-Lyte 148 Intravenous Solution
- Plasma-Lyte A Intravenous Solution
- Plenammine Intravenous Solution 15 %
- Potassium Chloride in Dextrose Intravenous Solution 20-5 MEQ/L-%, 40-5 MEQ/L-%
- Potassium Chloride in NaCl Intravenous Solution 20-0.45 MEQ/L-%, 20-0.9 MEQ/L-%, 40-0.9 MEQ/L-%
- Potassium Chloride Intravenous Solution 10 MEQ/100ML, 2 MEQ/ML (20 ML), 20 MEQ/100ML, 40 MEQ/100ML
- Premasol Intravenous Solution 10 %, 6 %
- Procalamine Intravenous Solution 3 %
- Prochlorperazine Edisylate Injection Solution 5 MG/ML
- Prograf Intravenous Solution 5 MG/ML
- Propranolol HCl Intravenous Solution 1 MG/ML
- Prosol Intravenous Solution 20 %
- RabAvert Intramuscular Suspension Reconstituted
- Rapamune Oral Solution 1 MG/ML
- Recombivax HB Injection Suspension 10 MCG/ML, 10 MCG/ML (1ML SYRINGE), 40 MCG/ML, 5 MCG/0.5ML
- Rifampin Intravenous Solution Reconstituted 600 MG
- Ringers Intravenous Solution
- Rituxan Intravenous Solution 100 MG/10ML, 500 MG/50ML
- SandIMMUNE Oral Capsule 100 MG, 25 MG
- SandIMMUNE Oral Solution 100 MG/ML
- Sensipar Oral Tablet 30 MG, 60 MG, 90 MG
- Sirolimus Oral Tablet 0.5 MG, 1 MG, 2 MG
- Sodium Chloride Intravenous Solution 0.45 %, 0.9 %, 3 %, 5 %
- Sodium Lactate Intravenous Solution 5 MEQ/ML
- Sulfamethoxazole-Trimethoprim Intravenous Solution 400-80 MG/5ML
- Synercid Intravenous Solution Reconstituted 150-350 MG
- Tacrolimus Oral Capsule 0.5 MG, 1 MG, 5 MG
- Teflaro Intravenous Solution Reconstituted 400 MG, 600 MG
- Tenivac Intramuscular Injectable 5-2 LFU

- Tetanus-Diphtheria Toxoids Td Intramuscular Suspension 2-2 LF/0.5ML
- Thiotepa Injection Solution Reconstituted 100 MG/ML 15 MG
- Thymoglobulin Intravenous Solution Reconstituted 25 MG
- Tigecycline Intravenous Solution Reconstituted 50 MG
- Toposar Intravenous Solution 1 GM/50ML
- Topotecan HCl Intravenous Solution Reconstituted 4 MG
- Torisel Intravenous Solution 25 MG/ML
- TPN Electrolytes Intravenous Solution
- Tranexamic Acid Intravenous Solution 1000 MG/10ML
- Travasol Intravenous Solution 10 %
- Treanda Intravenous Solution Reconstituted 100 MG, 25 MG
- Trexall Oral Tablet 10 MG, 15 MG, 5 MG, 7.5 MG
- Trisenox Intravenous Solution 12 MG/6ML
- TrophAmine Intravenous Solution 10 %
- Valproate Sodium Intravenous Solution
- Vancomycin HCl Intravenous Solution Reconstituted 10 GM, 1000 MG, 500 MG
- Varubi Oral Tablet 90 MG
- Verapamil HCl Intravenous Solution 2.5 MG/ML
- Vimpat Intravenous Solution 200 MG/20ML
- VinBLASTine Sulfate Intravenous Solution 1 MG/ML
- VinCRISTine Sulfate Intravenous Solution 1 MG/ML
- Vinorelbine Tartrate Intravenous Solution 50 MG/5ML
- Voriconazole Intravenous Solution Reconstituted 200 MG
- Xatmep Oral Solution 2.5 MG/ML
- Zanosar Intravenous Solution Reconstituted 1 GM
- Zerbaxa Intravenous Solution Reconstituted 1.5 (1-0.5) GM

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Index of Drugs/Alphabetical Listing

A

Abelcet.....	249
Abraxane	249
Abstral	64
Acetylcysteine	249
Acitretin	1
Actemra.....	2
Acyclovir Sodium.....	249
Adagen.....	3
Adapalene External Cream	212
Adapalene External Gel	212
Adcirca.....	4
Adefovir Dipivoxil.....	76
Adempas	5
Adrucil	249
Afinitor.....	6
Afinitor Disperz	6
Albuterol Sulfate	249
Aldurazyme.....	7
Alecensa.....	8
Alimta	249
Aliqopa.....	9
Alunbrig	11
AmBisome	249
Amikacin Sulfate	249
Aminosyn II	249
Aminosyn II/Electrolytes	249
Aminosyn/Electrolytes	249
Aminosyn-HBC	249
Aminosyn-PF.....	249
Aminosyn-RF.....	249
Amiodarone HCl	249
Amitriptyline HCl Oral	84
Amphotericin B	249
Ampicillin-Sulbactam Sodium	249
Ampyra.....	12
Apokyn Subcutaneous Solution Cartridge	13
Aprepitant	249
Aranesp (Albumin Free) Injection Solution 100 MCG/ML, 200 MCG/ML, 25 MCG/ML, 300 MCG/ML, 40 MCG/ML, 60 MCG/ML.....	58

Aranesp (Albumin Free) Injection Solution Prefilled Syringe	58
Arcalyst	14
Argatroban	249
Armodafinil	15
Arranon	249
Astagraf XL	249
Atgam	249
Aubagio	16
Auryxia	17
Austedo	18
Avastin	19
AzaCITIDine.....	249
Azactam	249
Azasan.....	249
AzaTHIOprine.....	249
AzaTHIOprine Sodium	249
Azithromycin	249
B	
Baraclude Oral Solution.....	76
Bavencio.....	21
Beleodaq	22
Benlysta	23
Benztropine Mesylate Injection.....	88
Benztropine Mesylate Oral	88
Betaseron Subcutaneous Kit	137
Bexarotene.....	24
BiCNU.....	249
Bleomycin Sulfate.....	249
Bortezomib	222
Bosulif.....	25
Briviact.....	26
Budesonide.....	249
Bupap Oral Tablet 50-300 MG.....	80
Buprenorphine HCl Sublingual Tablet Sublingual 2 MG, 8 MG.....	27
Busulfan.....	249
Butalbital-Acetaminophen Oral Tablet 50-325 MG	80
Butalbital-APAP-Caff-Cod Oral Capsule 50-325-40-30 MG	80
Butalbital-APAP-Caffeine Oral Tablet 50-325-40 MG	80

Butalbital-ASA-Caff-Codeine	80	Clinimix/Dextrose (4.25/10)	250
Butisol Sodium Oral Tablet 30 MG ...	95	Clinimix/Dextrose (4.25/20)	250
C		Clinimix/Dextrose (4.25/25)	250
Cabometyx	28	Clinimix/Dextrose (4.25/5)	250
Calcitonin (Salmon)	249	Clinimix/Dextrose (5/15)	250
Calcitriol.....	249	Clinimix/Dextrose (5/20)	250
Calquence	29	Clinimix/Dextrose (5/25)	250
Caprelsa	30	Clinisol SF	250
Carbaglu	31	Clofarabine	250
Carbinoxamine Maleate Oral Tablet 4		ClomiPRAMINE HCl Oral	84
MG	85	Cometriq (100 mg Daily Dose).....	37
CARBOplatin.....	249	Cometriq (140 mg Daily Dose).....	37
Carimune NF Intravenous Solution		Cometriq (60 mg Daily Dose).....	37
Reconstituted 6 GM.....	32	Corlanor.....	39
Carisoprodol Oral Tablet 350 MG.....	96	Cosentyx 300 Dose	40
Caspofungin Acetate	249	Cosentyx Sensoready 300 Dose	40
Cayston	34	Cotellic	41
CefOXitin Sodium	249	Cromolyn Sodium.....	250
CefTRIAXone Sodium	249	Cyclobenzaprine HCl Oral Tablet 10 MG,	
Cerezyme.....	249	5 MG	96
Chloramphenicol Sod Succinate	249	Cyclophosphamide	250
Chlordiazepoxide-Amitriptyline	84	CycloSPORINE	250
ChlorproMAZINE HCl Injection Solution		CycloSPORINE Modified.....	250
50 MG/2ML	90	Cyproheptadine HCl Oral.....	85
ChlorproMAZINE HCl Oral.....	90	Cyramza.....	250
ChlorproPAMIDE.....	97	Cystaran	42
Chlorzoxazone Oral Tablet 500 MG	96	Cytarabine.....	250
Chorionic Gonadotropin Intramuscular	73	Cytarabine (PF).....	250
Cidofovir	249	D	
Cimzia Prefilled.....	35	Dacarbazine	250
Cimzia Subcutaneous Kit 2 X 200 MG	35	DACTINomycin	250
Cinryze	36	Daliresp	43
Ciprofloxacin in D5W	250	DAPTOMycin	250
CISplatin	250	Darzalex Intravenous Solution 100	
Cladribine.....	250	MG/5ML	44
Clindamycin Phosphate	250	DAUNOrubicin HCl	250
Clinimix E/Dextrose (2.75/10)	250	Decitabine	250
Clinimix E/Dextrose (2.75/5)	250	Depo-Provera	250
Clinimix E/Dextrose (4.25/10)	250	Dexamethasone Sodium Phosphate ..	250
Clinimix E/Dextrose (4.25/25)	250	Dexrazoxane	250
Clinimix E/Dextrose (4.25/5)	250	Dextrose	250
Clinimix E/Dextrose (5/15)	250	Dextrose in Lactated Ringers	250
Clinimix E/Dextrose (5/20)	250	Dextrose-NaCl.....	250
Clinimix E/Dextrose (5/25)	250	Diclofenac Sodium Transdermal Gel...	45
Clinimix/Dextrose (2.75/5)	250	Digitek Oral Tablet 250 MCG	81

Digox Oral Tablet 250 MCG	81	Erleada	55
Digoxin Injection	81	Erwinaze Injection.....	56
Digoxin Oral Solution	81	Erythrocin Lactobionate.....	251
Digoxin Oral Tablet 250 MCG.....	81	Esbriet.....	57
Diltiazem HCl.....	250	Esomeprazole Sodium	251
Diphtheria-Tetanus Toxoids DT.....	251	Estradiol Oral	94
Dipyridamole Oral	89	Estradiol Transdermal.....	94
Disopyramide Phosphate Oral	81	Estropipate Oral Tablet 0.75 MG	94
Divigel Transdermal Gel 1 MG/GM	94	Etopophos.....	251
DOCEtaxel	251	Etoposide.....	251
Doxepin HCl Oral	84	Evamist	94
DOXOrubicin HCl.....	251	Exjade	59
DOXOrubicin HCl Liposomal.....	251	F	
Doxy 100.....	251	Fabrazyme	60
Dronabinol	251	Famotidine.....	251
E		Famotidine Premixed.....	251
Elaprase	46	Fareston	61
Elestrin	94	Farydak	62
Eligard	130	Faslodex	251
Elitek	251	FentaNYL.....	63
Emend	251	Firazyr	65
Empliciti	47	Firmagon	66
Emsam.....	48	Flebogamma DIF	251
Enbrel Subcutaneous Solution Prefilled Syringe.....	49	Fluconazole in Sodium Chloride.....	251
Enbrel Subcutaneous Solution Reconstituted	49	Fludarabine Phosphate	251
Enbrel SureClick Subcutaneous Solution Auto-Injector.....	49	Fluorouracil.....	251
Endari	51	Folotyln	251
Engerix-B.....	251	Forteo Subcutaneous Solution 600 MCG/2.4ML	67
Entecavir	76	FreAmine HBC.....	251
Entresto.....	52	G	
Envarsus XR.....	251	GamaSTAN S/D	251
Epclusa	77	Gammagard.....	251
Epirubicin HCl	251	Gammagard S/D Less IgA.....	251
Epogen Injection Solution 10000 UNIT/ML, 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML	58	Gammaplex	251
Eraxis.....	251	Ganciclovir Sodium	251
Erbitux Intravenous Solution 100 MG/50ML.....	53	Gattex	68
Ergoloid Mesylates Oral	83	Gemcitabine HCl	251
Erivedge	54	Gengraf	251
		Gilenya Oral Capsule 0.5 MG	69
		Gilotrif.....	70
		Glatiramer Acetate	38
		Gleostine Oral Capsule 10 MG, 100 MG, 40 MG	71
		GlyBURIDE Micronized.....	97

GlyBURIDE Oral	97	Intralipid.....	252
GlyBURIDE-MetFORMIN	97	Intrarosa	110
Gocovri.....	72	Intron A	111
Granisetron HCl.....	251	Ipratropium Bromide	252
GuanFACINE HCl ER	92	Ipratropium-Albuterol	252
GuanFACINE HCl Oral.....	92	Iressa	112
H		Irinotecan HCl	252
Halaven	75	Isolyte-S	252
Heparin (Porcine) in D5W.....	251	Istodax (Overfill)	113
Heparin Sod (Porcine) in D5W.....	251	Itraconazole Oral Capsule.....	190
Hepatamine	251	J	
Herceptin	78	Jakafi	114
Hexalen	79	Jevtana	252
Humira Pediatric Crohns Start		Juxtapid.....	115
Subcutaneous Prefilled Syringe Kit.	98	K	
Humira Pen Subcutaneous Pen-Injector		Kadcyla	116
Kit.....	98	Kalydeco.....	117
Humira Pen-CD/UC/HS Starter	98	Kanuma.....	118
Humira Pen-Ps/UV Starter	98	KCl in Dextrose-NaCl.....	252
Humira Subcutaneous Prefilled Syringe		KCl-Lactated Ringers-D5W.....	252
Kit.....	98	Kepivance.....	252
Hydroxyprogesterone Caproate		Ketorolac Tromethamine Injection	
Intramuscular Solution.....	100	Solution 15 MG/ML, 30 MG/ML	86
HydrOXYzine HCl Intramuscular	85	Ketorolac Tromethamine Intramuscular	
HydrOXYzine HCl Oral Syrup	85	Solution 60 MG/2ML	86
HydrOXYzine HCl Oral Tablet	85	Ketorolac Tromethamine Oral	86
HydrOXYzine Pamoate Oral	85	Keytruda Intravenous Solution	119
I		Kineret Subcutaneous Solution Prefilled	
Ibandronate Sodium.....	251	Syringe.....	121
Ibrance.....	101	Kisqali 200 Dose	122
Iclusig.....	102	Kisqali 400 Dose	122
IDArubicin HCl.....	251	Kisqali 600 Dose	122
IDHIFA.....	103	Kisqali Femara 200 Dose.....	122
Ifosfamide	251	Kisqali Femara 400 Dose.....	122
Imatinib Mesylate	104	Kisqali Femara 600 Dose.....	122
Imbruvica	105	Korlym.....	123
Imfinzi	106	Kuvan.....	124
Imipenem-Cilastatin.....	251	Kynamro Subcutaneous Solution	
Imipramine HCl Oral	84	Prefilled Syringe	125
Imipramine Pamoate.....	84	Kyprolis	252
Imogam Rabies-HT	251	L	
Increlex	107	Labetalol HCl	252
Indomethacin ER.....	86	Lactated Ringers.....	252
Indomethacin Oral	86	Lartruvo	126
Inlyta	109	Lenvima 10 MG Daily Dose	127

Lenvima 14 MG Daily Dose.....	127	Methocarbamol Injection Solution 1000 MG/10ML.....	96
Lenvima 18 MG Daily Dose.....	127	Methocarbamol Oral.....	96
Lenvima 20 MG Daily Dose.....	127	Methotrexate.....	252
Lenvima 24 MG Daily Dose.....	127	Methotrexate Sodium.....	252
Lenvima 8 MG Daily Dose.....	127	Methotrexate Sodium (PF).....	252
Letairis.....	128	Methyldopa Oral.....	92
Leucovorin Calcium.....	252	Methyldopa-Hydrochlorothiazide.....	92
Leukine Intravenous.....	129	Methyldopate HCl.....	92
Leuprolide Acetate Injection.....	130	MethylPREDNISolone Sodium Succ.....	252
LevETIRAcetam.....	252	Metoprolol Tartrate.....	252
LevETIRAcetam in NaCl.....	252	MetroNIDAZOLE in NaCl.....	252
Levofloxacin.....	252	Miacalcin.....	252
Levofloxacin in D5W.....	252	Miglustat.....	240
Levoleucovorin Calcium.....	252	Mircera Injection Solution Prefilled Syringe 100 MCG/0.3ML, 50 MCG/0.3ML, 75 MCG/0.3ML.....	58
LEVOleucovorin Calcium.....	252	MitoMYcin.....	252
Lidocaine External Patch 5 %.....	131	Mitoxantrone HCl.....	252
Linezolid Intravenous Solution 600 MG/300ML.....	132	Modafinil.....	15
Linezolid Oral Tablet.....	132	Moxifloxacin HCl in NaCl.....	252
Lonsurf.....	133	Mozobil.....	136
Lumizyme.....	252	Mustargen.....	252
Lupron Depot (1-Month).....	130	Mycamine.....	252
Lupron Depot (3-Month).....	130	Mycophenolate Mofetil.....	252
Lupron Depot (4-Month).....	130	Mycophenolate Mofetil HCl.....	252
Lupron Depot (6-Month).....	130	Mycophenolate Sodium.....	252
Lupron Depot-Ped (1-Month) Intramuscular Kit 11.25 MG, 15 MG	130	Mylotarg Intravenous Solution Reconstituted 4.5 MG.....	138
Lupron Depot-Ped (3-Month) Intramuscular Kit 30 MG (Ped)....	130	Mytesi.....	139
Lynparza.....	134	N	
M		Naglazyme.....	140
Mavyret.....	77	Natpara.....	141
Megestrol Acetate Oral Suspension 40 MG/ML, 625 MG/5ML.....	87	Nebupent.....	253
Megestrol Acetate Oral Tablet.....	87	NephrAmine.....	253
Mekinist.....	135	Nerlynx.....	142
Melphalan HCl.....	252	Neupogen Injection Solution 300 MCG/ML, 480 MCG/1.6ML.....	143
Menest Oral Tablet 0.3 MG, 0.625 MG, 1.25 MG.....	94	Neupogen Injection Solution Prefilled Syringe.....	143
Meperidine HCl Injection Solution 100 MG/ML, 25 MG/ML.....	80	NexAVAR.....	145
Meprobamate.....	91	NIFEdipine Oral.....	92
Meropenem.....	252	Ninlaro.....	146
Mesna.....	252	Nitroglycerin.....	253
		Norditropin FlexPro.....	74

Norethindrone-Eth Estradiol.....	94	Pentazocine-Naloxone HCl	80
Normosol-M in D5W.....	253	Perjeta.....	165
Normosol-R in D5W	253	Perphenazine Oral.....	90
Normosol-R pH 7.4.....	253	Perphenazine-Amitriptyline	84
Northera	147	Phenadoz Rectal Suppository 25 MG..	85
Noxafil Oral.....	148	PHENobarbital Oral Elixir.....	82
Nucala	149	PHENobarbital Oral Tablet.....	82
Nuedexta.....	150	Piperacillin Sod-Tazobactam So.....	253
Nulojix	151	Plasma-Lyte 148	253
Nuplazid Oral Tablet 17 MG	152	Plasma-Lyte A	253
Nutrilipid.....	253	Plenamaine.....	253
O		Pomalyst.....	166
Octreotide Acetate Injection Solution		Potassium Chloride	253
100 MCG/ML, 1000 MCG/ML, 200		Potassium Chloride in Dextrose.....	253
MCG/ML, 50 MCG/ML, 500 MCG/ML		Potassium Chloride in NaCl.....	253
.....	182	Praluent Subcutaneous Solution Pen-	
Odomzo.....	153	Injector	162
Ofev.....	169	Premasol.....	253
Ondansetron	253	Procalamine	253
Ondansetron HCl	253	Prochlorperazine Edisylate	253
Opdivo Intravenous Solution 100		Procrit	58
MG/10ML, 40 MG/4ML.....	154	Prograf	253
Opsumit	156	Prolastin-C Intravenous Solution	
Orencia ClickJect.....	157	Reconstituted 1000 MG	10
Orencia Intravenous	157	Proleukin	167
Orencia Subcutaneous Solution Prefilled		Promacta	168
Syringe.....	157	Promethazine HCl Injection.....	85
Orkambi Oral Tablet.....	158	Promethazine HCl Oral Syrup	85
Orphenadrine Citrate ER.....	96	Promethazine HCl Oral Tablet.....	85
Orphenadrine Citrate Injection.....	96	Promethazine HCl Rectal Suppository	
Osphena	159	12.5 MG	85
Otrexup Subcutaneous Solution Auto-		Promethegan Rectal Suppository 12.5	
Injector 10 MG/0.4ML, 12.5		MG	85
MG/0.4ML, 15 MG/0.4ML, 20		Propranolol HCl	253
MG/0.4ML, 22.5 MG/0.4ML, 25		Prosol.....	253
MG/0.4ML.....	160	Pulmozyme	170
Oxaliplatin	253	Q	
Oxandrolone Oral	161	QuiNINE Sulfate Oral	171
P		R	
PACLitaxel	253	RabAvert.....	253
Pamidronate Disodium	253	Radicava.....	172
Pantoprazole Sodium	253	Ranexa	173
Paricalcitol	253	Rapamune	253
Pegasys ProClick	164	Rasuvo Subcutaneous Solution Auto-	
Pegasys Subcutaneous Solution.....	164	Injector 10 MG/0.2ML, 12.5	

MG/0.25ML, 15 MG/0.3ML, 17.5 MG/0.35ML, 20 MG/0.4ML, 22.5 MG/0.45ML, 25 MG/0.5ML, 30 MG/0.6ML, 7.5 MG/0.15ML.....	174	Sylatron Subcutaneous Kit 200 MCG, 300 MCG, 600 MCG.....	196
Recombivax HB.....	253	Symdeko.....	197
Regranex	175	SymlinPen 120 Subcutaneous Solution Pen-Injector	198
Repatha.....	162	SymlinPen 60 Subcutaneous Solution Pen-Injector	198
Repatha Pushtronex System.....	162	Synagis.....	199
Repatha SureClick	162	Synarel	201
Revlimid	176	Synercid	253
Rifampin	253	Synribo	202
Riluzole.....	177	T	
Ringers.....	253	Tacrolimus	253
Rituxan	253	Tafinlar.....	204
Rubraca.....	178	Tagrisso	205
Rydapt	179	Tarceva.....	206
S		Tasigna.....	207
Sabril Oral Tablet	180	Tazarotene External.....	208
Samsca	181	Tazorac External Cream 0.05 %.....	208
SandIMMUNE	253	Tazorac External Gel.....	208
SandoSTATIN LAR Depot	182	Tecentriq	209
Sensipar	253	Teflaro.....	253
Signifor	183	Tenivac.....	253
Sildenafil Citrate Intravenous	184	Testosterone Cypionate Intramuscular Solution 100 MG/ML, 200 MG/ML.....	210
Sildenafil Citrate Oral Tablet 20 MG	184	Testosterone Enanthate Intramuscular Solution.....	210
Simponi Aria.....	185	Testosterone Transdermal Gel 10 MG/ACT (2%), 12.5 MG/ACT (1%), 50 MG/5GM (1%).....	210
Simponi Subcutaneous Solution Auto- Injector	185	Testosterone Transdermal Solution...	210
Simponi Subcutaneous Solution Prefilled Syringe.....	185	Tetanus-Diphtheria Toxoids Td	254
Sirolimus.....	253	Tetrabenazine.....	211
Sodium Chloride.....	253	Thalomid	176
Sodium Lactate.....	253	Thioridazine HCl Oral	90
Soltamox	187	Thiotepa	254
Somatuline Depot	188	Thymoglobulin.....	254
Somavert	189	Tigecycline.....	254
Sprycel.....	191	Tobi Podhaler	108
Stelara Intravenous.....	192	Tobramycin Inhalation	108
Stelara Subcutaneous Solution 45 MG/0.5ML.....	192	Toposar	254
Stelara Subcutaneous Solution Prefilled Syringe.....	192	Topotecan HCl.....	254
Stivarga.....	194	Torisel	254
Sulfamethoxazole-Trimethoprim	253	TPN Electrolytes	254
Sutent.....	195	Tracleer	213

Tranexamic Acid.....	254	Vyxeos Intravenous Suspension	
Travasol.....	254	Reconstituted 44-100 MG.....	228
Treanda.....	254	X	
Trelstar Mixject	214	Xalkori.....	229
Tretinoin External Cream	212	Xatmep.....	254
Tretinoin External Gel 0.01 %, 0.025 %		Xeljanz	230
.....	212	Xeljanz XR	230
Trexall.....	254	Xgeva	231
Trientine HCl	203	Xolair.....	232
Trihexyphenidyl HCl.....	88	Xtandi	233
Trimipramine Maleate Oral.....	84	Xuriden.....	234
Trisenox.....	254	Xyrem.....	235
TrophAmine	254	Y	
Tykerb.....	215	Yervoy Intravenous Solution 50	
Tymlos	216	MG/10ML.....	236
Tysabri	217	Yondelis.....	237
U		Yonsa	238
Uptravi	218	Z	
V		Zaleplon Oral Capsule 10 MG, 5 MG ..	95
Valchlor.....	219	Zaltrap Intravenous Solution 100	
Valproate Sodium	254	MG/4ML.....	239
Vancomycin HCl.....	254	Zanosar	254
VariZIG Intramuscular Solution.....	220	Zarxio.....	143
Varubi	254	Zavesca	240
Vectibix Intravenous Solution 100		Zejula.....	241
MG/5ML.....	221	Zelboraf.....	242
Velcade Injection.....	222	Zemaira	10
Vemlidy.....	76	Zepatier.....	77
Venclexta.....	223	Zerbaxa	254
Venclexta Starting Pack.....	223	Zoledronic Acid Intravenous Concentrate	
Ventavis.....	224	243
Verapamil HCl	254	Zoledronic Acid Intravenous Solution 5	
Verzenio	225	MG/100ML.....	243
Vigabatrin.....	180	Zolinza	244
Vimpat	254	Zolpidem Tartrate ER	95
VinBLAStine Sulfate	254	Zolpidem Tartrate Oral Tablet 10 MG, 5	
VinCRIStine Sulfate	254	MG	95
Vinorelbine Tartrate	254	Zolpidem Tartrate Sublingual	95
Voriconazole.....	254	Zortress	245
Votrient	226	Zydelig	246
Vpriv	227	Zykadia.....	247
		Zytiga	248